




Rec'd PCT/PTO 18 AUG 2005
PCT/GB2004/000695
1 546005

INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ



I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

R. Mahoney

Dated 16 March 2004

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Best Available Copy

Patents Form 1/77

Patents Act 1977
Rule 16)

THE PATENT OFFICE

25 FEB 2003

NEWPORT

The
Patent
Office

25 FEB 2003 E782610-1 C76921
P01/2000 0300-0304194.4

Request for grant of a patent

See the notes on the back of this form. You can also get
an explanatory leaflet from the Patent Office to help
you fill in this form.

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

Your reference

100994

Patent application number

(The Patent Office will fill in this part)

25 FEB 2003

0304194.4

Full name, address and postcode of the or of
each applicant (underline all surnames)

AstraZeneca AB
SE-151 85 Sodertalje
Sweden

Patents ADP number (if you know it)

If the applicant is a corporate body, give the
country/state of its incorporation

Sweden

7954274002

Title of the invention

CHEMICAL COMPOUNDS

Name of your agent (if you have one)

Lucy Clare Padget

"Address for service" in the United Kingdom
to which all correspondence should be sent
(including the postcode)

AstraZeneca UK Limited
Global Intellectual Property
Mereside, Alderley Park
Macclesfield
Cheshire SK10 4TG

Patents ADP number (if you know it)

7822471002

If you are declaring priority from one or more
earlier patent applications, give the country
and the date of filing of the or of each of these
earlier applications and (if you know it) the or
each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

If this application is divided or otherwise
derived from an earlier UK application,
give the number and the filing date of
the earlier application

Number of earlier application

Date of filing
(day / month / year)

Is a statement of inventorship and of right
to grant of a patent required in support of
this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an
applicant, or
- c) any named applicant is a corporate body.

See note (d))

Patents Form 1/77

- Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 55

Claim(s) 06

Abstract 01

Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date

Authorised Signatory

24/02/2003

12. Name and daytime telephone number of person to contact in the United Kingdom

Diana Fletcher - 01625 515907

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

CHEMICAL COMPOUNDS

This invention relates to benzothiazepine and benzothiepine derivatives, or pharmaceutically acceptable salts, solvates, solvates of such salts and prodrugs thereof. These benzothiazepines and benzothiepies possess ileal bile acid transport (IBAT) inhibitory activity and accordingly have value in the treatment of disease states associated with hyperlipidaemic conditions and they are useful in methods of treatment of a warm-blooded animal, such as man. The invention also relates to processes for the manufacture of said benzothiazepine and benzothiepine derivatives, to pharmaceutical compositions containing them and to their use in the manufacture of medicaments to inhibit IBAT in a warm-blooded animal, such as man.

It is well-known that hyperlipidaemic conditions associated with elevated concentrations of total cholesterol and low-density lipoprotein cholesterol are major risk factors for cardiovascular atherosclerotic disease (for instance "Coronary Heart Disease: Reducing the Risk; a Worldwide View" Assman G., Carmena R. Cullen P. *et al*; Circulation 1999, 100, 1930-1938 and "Diabetes and Cardiovascular Disease: A Statement for Healthcare Professionals from the American Heart Association" Grundy S, Benjamin I., Burke G., *et al*; Circulation, 1999, 100, 1134-46). Interfering with the circulation of bile acids within the lumen of the intestinal tracts is found to reduce the level of cholesterol. Previous established therapies to reduce the concentration of cholesterol involve, for instance, treatment with HMG-CoA reductase inhibitors, preferably statins such as simvastatin and fluvastatin, or treatment with bile acid binders, such as resins. Frequently used bile acid binders are for instance cholestyramine and cholestipol. One recently proposed therapy ("Bile Acids and Lipoprotein Metabolism: a Renaissance for Bile Acids in the Post Statin Era" Angelin B, Eriksson M, Rudling M; Current Opinion on Lipidology, 1999, 10, 269-74) involved the treatment with substances with an IBAT inhibitory effect.


Re-absorption of bile acid from the gastro-intestinal tract is a normal physiological process which mainly takes place in the ileum by the IBAT mechanism. Inhibitors of IBAT can be used in the treatment of hypercholesterolaemia (see for instance "Interaction of bile acids and cholesterol with nonsystemic agents having hypocholesterolaemic properties", Biochimica et Biophysica Acta, 1210 (1994) 255- 287). Thus, suitable compounds having such inhibitory IBAT activity are also useful in the treatment of hyperlipidaemic conditions. Substituted compounds possessing such IBAT inhibitory activity have been described, see for instance

hypolipidaemic compounds described in WO 93/16055, WO 94/18183, WO 94/18184, WO 96/05188, WO 96/08484, WO 96/16051, WO 97/33882, WO 98/38182, WO 99/35135, WO 98/40375, WO 99/35153, WO 99/64409, WO 99/64410, WO 00/01687, WO 00/47568, WO 00/61568, WO 01/68906, DE 19825804, WO 00/38725, WO 00/38726, WO 00/38727, WO 5 00/38728, WO 00/38729, WO 01/68906 and EP 0 864 582.

A further aspect of this invention relates to the use of the compounds of the invention in the treatment of dyslipidemic conditions and disorders such as hyperlipidaemia, hypertriglyceridemia, hyperbetalipoproteinemia (high LDL), hyperprebetalipoproteinemia (high VLDL), hyperchylomicronemia, hypolipoproteinemia, hypercholesterolemia, 10 hyperlipoproteinemia and hypoalphalipoproteinemia (low HDL). In addition, these compounds are expected to be useful for the prevention and treatment of different clinical conditions such as atherosclerosis, arteriosclerosis, arrhythmia, hyper-thrombotic conditions, vascular dysfunction, endothelial dysfunction, heart failure, coronary heart diseases, cardiovascular diseases, myocardial infarction, angina pectoris, peripheral vascular diseases, inflammation of 15 cardiovascular tissues such as heart, valves, vasculature, arteries and veins, aneurisms, stenosis, restenosis, vascular plaques, vascular fatty streaks, leukocytes, monocytes and/or macrophage infiltration, intimal thickening, medial thinning, infectious and surgical trauma and vascular thrombosis, stroke and transient ischaemic attacks.

The present invention is based on the discovery that certain benzothiazepine and 20 benzothiepine compounds surprisingly inhibit IBAT. Such properties are expected to be of value in the treatment of disease states associated with hyperlipidaemic conditions.

Accordingly, the present invention provides a compound of formula (I):



25 wherein

M^1 is $-CH_2-$ or $-NR^{21}-$;


M^2 is $-CR^{22}R^{23}-$ or $-NR^{24}-$; provided that if M^1 is $-NR^{21}-$, M^2 is $-CR^{22}R^{23}-$;

One of R^1 and R^2 are selected from hydrogen, C_{1-6} alkyl or C_{2-6} alkenyl and the other is selected from C_{1-6} alkyl or C_{2-6} alkenyl;

R^3 is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, $N-(C_{1-6}alkyl)amino$, $N,N-(C_{1-6}alkyl)_2amino$, $C_{1-6}alkanoylamino$, $N-(C_{1-6}alkyl)carbamoyl$, $N,N-(C_{1-6}alkyl)_2carbamoyl$, $C_{1-6}alkylS(O)_a$ wherein a is 0 to 2, $C_{1-6}alkoxycarbonyl$, $N-(C_{1-6}alkyl)sulphamoyl$ and $N,N-(C_{1-6}alkyl)_2sulphamoyl$;

v is 0-5;

10 one of R^5 and R^6 is a group of formula (IA):



(IA)

R^4 and R^7 and the other of R^5 and R^6 are independently selected from hydrogen, halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, C_{1-4} alkanoyl, C_{1-4} alkanoyloxy, $N-(C_{1-4}alkyl)amino$, $N,N-(C_{1-4}alkyl)_2amino$, $C_{1-4}alkanoylamino$, $N-(C_{1-4}alkyl)carbamoyl$, $N,N-(C_{1-4}alkyl)_2carbamoyl$, $C_{1-4}alkylS(O)_a$ wherein a is 0 to 2, $C_{1-4}alkoxycarbonyl$, $N-(C_{1-4}alkyl)sulphamoyl$ and $N,N-(C_{1-4}alkyl)_2sulphamoyl$; wherein R^4 and R^7 and the other of R^5 and R^6 may be optionally substituted on carbon by one or more R^{25} ;

20 Z is $-O-$, $-N(R^a)-$, $-S(O)_b-$ or $-CH(R^a)-$; wherein R^a is hydrogen or C_{1-6} alkyl and b is 0-2;

R^8 is hydrogen, C_{1-4} alkyl, carbocyclyl or heterocyclyl; wherein R^8 may be optionally substituted on carbon by one or more substituents selected from R^{26} ; and wherein if said heterocyclyl contains an $-NH-$ group, that nitrogen may be optionally substituted by a group selected from R^{27} ;

R^9 is hydrogen or C_{1-4} alkyl;

R^{10} and R^{11} are independently selected from hydrogen, C_{1-4} alkyl, carbocyclyl or heterocyclyl; or R^{10} and R^{11} together form C_{2-6} alkylene; wherein R^{10} and R^{11} or R^{10} and R^{11} together may be independently optionally substituted on carbon by one or more substituents

selected from R^{28} ; and wherein if said heterocyclyl contains an -NH- moiety, that nitrogen may be optionally substituted by one or more R^{29} ;

R^{12} is hydrogen, C_{1-4} alkyl, carbocyclyl or heterocyclyl; wherein R^{12} may be optionally substituted on carbon by one or more substituents selected from R^{30} ; and wherein if said

5 heterocyclyl contains an -NH- moiety, that nitrogen may be optionally substituted by one or more R^{31} ;

R^{13} is hydrogen, halo, nitro, cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, hydroxyaminocarbonyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy, C_{1-10} alkoxycarbonyl, C_{1-10} alkanoyl, C_{1-10} alkanoyloxy, N -(C_{1-10} alkyl)amino, N,N -(C_{1-10} alkyl)₂amino,

10 N,N,N -(C_{1-10} alkyl)₃ammonio, C_{1-10} alkanoylamino, N -(C_{1-10} alkyl)carbamoyl,

N,N -(C_{1-10} alkyl)₂carbamoyl, C_{1-10} alkylS(O)_a wherein a is 0 to 2, N -(C_{1-10} alkyl)sulphamoyl,


N,N -(C_{1-10} alkyl)₂sulphamoyl, N -(C_{1-10} alkyl)sulphamoylamino,

N,N -(C_{1-10} alkyl)₂sulphamoylamino, C_{1-10} alkoxycarbonylamino, carbocyclyl,

carbocyclyl C_{1-10} alkyl, heterocyclic group, heterocyclyl C_{1-10} alkyl,

15 carbocyclyl-(C_{1-10} alkylene)_e- R^{32} -(C_{1-10} alkylene)_f or

heterocyclyl-(C_{1-10} alkylene)_g- R^{33} -(C_{1-10} alkylene)_h; wherein R^{13} may be optionally substituted on carbon by one or more substituents selected from R^{36} ; and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally substituted by a group selected from R^{37} ; or R^{13} is a group of formula (IB):



20

(IB)

wherein:

X is -N(R^{38})-, -N(R^{38})C(O)-, -O-, and -S(O)_a-; wherein a is 0-2 and R^{38} is hydrogen or C_{1-4} alkyl;

25 R^{14} is hydrogen or C_{1-4} alkyl;

R^{15} and R^{16} are independently selected from hydrogen, halo, nitro, cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkanoylamino, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkylS(O)_a wherein a is 0 to 2,


30 C_{1-6} alkoxycarbonyl, N -(C_{1-6} alkyl)sulphamoyl, N,N -(C_{1-6} alkyl)₂sulphamoyl, carbocyclyl or

heterocyclic group; wherein R^{15} and R^{16} may be independently optionally substituted on carbon by one or more substituents selected from R^{41} ; and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally substituted by a group selected from R^{42} ;

R^{17} is selected from hydrogen, halo, nitro, cyano, hydroxy, amino, carbamoyl,

5 mercapto, sulphamoyl, hydroxyaminocarbonyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy, C_{1-10} alkanoyl, C_{1-10} alkanoyloxy, N -(C_{1-10} alkyl)amino, N,N -(C_{1-10} alkyl) $_2$ amino, C_{1-10} alkanoylamino, N -(C_{1-10} alkyl)carbamoyl, C_{1-10} alkoxycarbonyl, N,N -(C_{1-10} alkyl) $_2$ carbamoyl, C_{1-10} alkylS(O) $_a$ wherein a is 0 to 2, N -(C_{1-10} alkyl)sulphamoyl, N,N -(C_{1-10} alkyl) $_2$ sulphamoyl, N -(C_{1-10} alkyl)sulphamoylamino, N,N -(C_{1-10} alkyl) $_2$ sulphamoylamino, carbocyclyl,

10 carbocyclyl C_{1-10} alkyl, heterocyclic group, heterocyclyl C_{1-10} alkyl, carbocyclyl-(C_{1-10} alkylene) $_e$ - R^{43} -(C_{1-10} alkylene) $_f$ or heterocyclyl-(C_{1-10} alkylene) $_g$ - R^{44} -(C_{1-10} alkylene) $_h$; wherein R^{17} may be optionally substituted on carbon by one or more substituents selected from R^{47} ; and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally substituted by a group selected from R^{48} ; or R^{17} is a group of formula (IC):



(IC)

wherein:

R^{18} is selected from hydrogen or C_{1-4} alkyl;

20 R^{19} is selected from hydrogen, halo, nitro, cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkylS(O) $_a$ wherein a is 0 to 2, C_{1-6} alkoxycarbonyl, N -(C_{1-6} alkyl)sulphamoyl, N,N -(C_{1-6} alkyl) $_2$ sulphamoyl, carbocyclyl or

25 heterocyclic group; where R^{19} may be independently optionally substituted on carbon by one or more substituents selected from R^{51} ; and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally substituted by a group selected from R^{52} ;

R^{20} is selected from halo, nitro, cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, hydroxyaminocarbonyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy,

30 C_{1-10} alkoxycarbonyl, C_{1-10} alkanoyl, C_{1-10} alkanoyloxy, N -(C_{1-10} alkyl)amino,

- N,N*-(C₁₋₁₀alkyl)₂amino, *N,N,N*-(C₁₋₁₀alkyl)₃ammonio, C₁₋₁₀alkanoylamino, *N*-(C₁₋₁₀alkyl)carbamoyl, *N,N*-(C₁₋₁₀alkyl)₂carbamoyl, C₁₋₁₀alkylS(O)_a wherein a is 0 to 2, *N*-(C₁₋₁₀alkyl)sulphamoyl, *N,N*-(C₁₋₁₀alkyl)₂sulphamoyl, *N*-(C₁₋₁₀alkyl)sulphamoylamino, *N,N*-(C₁₋₁₀alkyl)₂sulphamoylamino, C₁₋₁₀alkoxycarbonylamino, carbocyclyl,
- 5 carbocyclylC₁₋₁₀alkyl, heterocyclic group, heterocyclylC₁₋₁₀alkyl, carbocyclyl-(C₁₋₁₀alkylene)_e-R⁵³-(C₁₋₁₀alkylene)_f or heterocyclyl-(C₁₋₁₀alkylene)_g-R⁵⁴-(C₁₋₁₀alkylene)_h; wherein R²⁰ may be independently optionally substituted on carbon by one or more R⁵⁷; and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally substituted by a group selected from R⁵⁸;
- 10 **p** is 1-3; wherein the values of R¹⁵ may be the same or different; **q** is 0-1; **r** is 0-3; wherein the values of R¹⁶ may be the same or different; **m** is 0-2; wherein the values of R¹² may be the same or different; **n** is 1-2; wherein the values of R⁸ may be the same or different;
- 15 **z** is 0-3; wherein the values of R¹⁹ may be the same or different; R²¹ is selected from hydrogen or C₁₋₆alkyl; R²² and R²³ are independently selected from hydrogen, hydroxy, amino, mercapto, C₁₋₆alkyl, C₁₋₆alkoxy, *N*-(C₁₋₆alkyl)amino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkylS(O)_a wherein a is 0 to 2;
- 20 R²⁴ is selected from hydrogen, hydroxy, C₁₋₆alkyl, C₁₋₄alkoxy and C₁₋₆alkanoyloxy; R²⁵ is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₁₋₄alkoxy, C₁₋₄alkanoyl, C₁₋₄alkanoyloxy, *N*-(C₁₋₄alkyl)amino, *N,N*-(C₁₋₄alkyl)₂amino, C₁₋₄alkanoylamino, *N*-(C₁₋₄alkyl)carbamoyl, *N,N*-(C₁₋₄alkyl)₂carbamoyl, C₁₋₄alkylS(O)_a wherein a is 0 to 2, C₁₋₄alkoxycarbonyl,
- 25 *N*-(C₁₋₄alkyl)sulphamoyl and *N,N*-(C₁₋₄alkyl)₂sulphamoyl; wherein R²⁵, may be independently optionally substituted on carbon by one or more R⁶⁷;
- R²⁶, R²⁸, R³⁰, R³⁶, R⁴¹, R⁴⁷, R⁵¹ and R⁵⁷ are independently selected from halo, nitro, cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, hydroxyaminocarbonyl, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, C₁₋₁₀alkoxy, C₁₋₁₀alkanoyl, C₁₋₁₀alkanoyloxy, C₁₋₁₀alkoxycarbonyl,
- 30 *N*-(C₁₋₁₀alkyl)amino, *N,N*-(C₁₋₁₀alkyl)₂amino, *N,N,N*-(C₁₋₁₀alkyl)₃ammonio, C₁₋₁₀alkanoylamino, *N*-(C₁₋₁₀alkyl)carbamoyl, *N,N*-(C₁₋₁₀alkyl)₂carbamoyl, C₁₋₁₀alkylS(O)_a wherein a is 0 to 2, *N*-(C₁₋₁₀alkyl)sulphamoyl, *N,N*-(C₁₋₁₀alkyl)₂sulphamoyl, *N*-(C₁₋₁₀alkyl)sulphamoylamino,

N,N-(C₁₋₁₀alkyl)₂sulphamoylamino, C₁₋₁₀alkoxycarbonylamino, carbocyclyl, carbocyclylC₁₋₁₀alkyl, heterocyclic group, heterocyclylC₁₋₁₀alkyl, carbocyclyl-(C₁₋₁₀alkylene)_e-R⁵⁹-(C₁₋₁₀alkylene)_f or heterocyclyl-(C₁₋₁₀alkylene)_g-R⁶⁰-(C₁₋₁₀alkylene)_h; wherein R²⁶, R²⁸, R³⁰, R³⁶, R⁴¹, R⁴⁷, R⁵¹ and 5 R⁵⁷ may be independently optionally substituted on carbon by one or more R⁶³; and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally substituted by a group selected from R⁶⁴;

R²⁷, R²⁹, R³¹, R³⁷, R⁴², R⁴⁸, R⁵², R⁵⁸ and R⁶⁴ are independently selected from C₁₋₆alkyl, C₁₋₆alkanoyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl, 10 *N,N*-(C₁₋₆alkyl)₂sulphamoyl, C₁₋₆alkoxycarbonyl, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, benzyl, phenethyl, benzoyl, phenylsulphonyl and phenyl;

R³², R³³, R⁴³, R⁴⁴, R⁵³, R⁵⁴, R⁵⁹ and R⁶⁰ are independently selected from -O-, -NR⁶⁵-, -S(O)_x-, -NR⁶⁵C(O)NR⁶⁶-, -NR⁶⁵C(S)NR⁶⁶-, -OC(O)N=C-, -NR⁶⁵C(O)- or -C(O)NR⁶⁵-; wherein R⁶⁵ and R⁶⁶ are independently selected from hydrogen or C₁₋₆alkyl, and x is 0-2;

15 R⁶³ and R⁶⁷ are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, methyl, ethyl, methoxy, ethoxy, vinyl, allyl, ethynyl, methoxycarbonyl, formyl, acetyl, formamido, acetylamino, acetoxymethyl, methylamino, dimethylamino, *N*-methylcarbamoyl, *N,N*-dimethylcarbamoyl, methylthio, methylsulphinyl, mesyl, *N*-methylsulphamoyl and

20 *N,N*-dimethylsulphamoyl; and

e, f, g and h are independently selected from 0-2;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

In this specification the term "alkyl" includes both straight and branched chain alkyl groups but references to individual alkyl groups such as "propyl" are specific for the straight 25 chain version only. For example, "C₁₋₁₀alkyl" and "C₁₋₆alkyl" includes C₁₋₄alkyl, C₁₋₃alkyl, propyl, isopropyl and *t*-butyl. However, references to individual alkyl groups such as 'propyl' are specific for the straight chained version only and references to individual branched chain alkyl groups such as 'isopropyl' are specific for the branched chain version only. A similar convention applies to other radicals, for example "phenylC₁₋₆alkyl" would include 30 phenylC₁₋₆alkyl, benzyl, 1-phenylethyl and 2-phenylethyl. The term "halo" refers to fluoro, chloro, bromo and iodo.

Where optional substituents are chosen from "one or more" groups it is to be understood that this definition includes all substituents being chosen from one of the specified groups or the substituents being chosen from two or more of the specified groups.

A "heterocyclyl" is a saturated, partially saturated or unsaturated, mono or bicyclic ring
5 containing 3-12 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a -CH₂- group can optionally be replaced by a -C(O)- or a ring sulphur atom may be optionally oxidised to form the S-oxides. Preferably a "heterocyclyl" is a saturated, partially saturated or unsaturated, mono or bicyclic ring containing 5 or 6 atoms of which at least one atom is chosen from
10 nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a -CH₂- group can optionally be replaced by a -C(O)- or a ring sulphur atom may be optionally oxidised to form S-oxide(s). Examples and suitable values of the term "heterocyclyl" are thiazolidinyl, pyrrolidinyl, pyrrolinyl, 2-pyrrolidinonyl, 2,5-dioxopyrrolidinyl, 2-benzoxazolinonyl, 1,1-dioxotetrahydrothienyl, 2,4-dioxoimidazolidinyl,
15 2-oxo-1,3,4-(4-triazolinyl), 2-oxazolidinonyl, 5,6-dihydrouracilyl, 1,3-benzodioxolyl, 1,2,4-oxadiazolyl, 2-azabicyclo[2.2.1]heptyl, 4-thiazolidonyl, morpholino, 2-oxotetrahydrofuranyl, tetrahydrofuranyl, 2,3-dihydrobenzofuranyl, benzothienyl, tetrahydropyranyl, piperidyl, 1-oxo-1,3-dihydroisoindolyl, piperazinyl, thiomorpholino, 1,1-dioxothiomorpholino, tetrahydropyranyl, 1,3-dioxolanyl, homopiperazinyl, thienyl,
20 isoxazolyl, imidazolyl, pyrrolyl, thiadiazolyl, isothiazolyl, 1,2,4-triazolyl, 1,3,4-triazolyl, pyranyl, indolyl, pyrimidyl, thiazolyl, pyrazinyl, pyridazinyl, pyridyl, 4-pyridonyl, quinolyl and 1-isoquinolonyl.

A "heterocyclic group" is a saturated, partially saturated or unsaturated, mono or bicyclic ring containing 3-12 atoms of which at least one atom is chosen from nitrogen, sulphur
25 or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a -CH₂- group can optionally be replaced by a -C(O)- or a ring sulphur atom may be optionally oxidised to form the S-oxides. A "heterocyclic group" is not tetrazolyl. Preferably a "heterocyclic group" is a saturated, partially saturated or unsaturated, monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 9 or 10 atoms of which at least one atom
30 is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a -CH₂- group can optionally be replaced by a -C(O)- or a ring sulphur atom may be optionally oxidised to form S-oxide(s). Examples and suitable values of

the term "heterocyclic group" are thiazolidinyl, pyrrolidinyl, pyrrolinyl, 2-pyrrolidonyl, 2,5-dioxopyrrolidinyl, 2-benzoxazolinonyl, 1,1-dioxotetrahydrothienyl, 2,4-dioxoimidazolidinyl, 2-oxo-1,3,4-(4-triazolinyl), 2-oxazolidinonyl, 5,6-dihydrouracilyl, 1,3-benzodioxolyl, 1,2,4-oxadiazolyl, 2-azabicyclo[2.2.1]heptyl, 4-thiazolidonyl, morpholino, 5 2-oxotetrahydrofuranyl, tetrahydrofuranyl, 2,3-dihydrobenzofuranyl, benzothieryl, tetrahydropyranyl, piperidyl, 1-oxo-1,3-dihydroisoindolyl, piperazinyl, thiomorpholino, 1,1-dioxothiormorpholino, tetrahydropyranyl, 1,3-dioxolanyl, homopiperazinyl, thienyl, isoxazolyl, imidazolyl, pyrrolyl, thiadiazolyl, isothiazolyl, 1,2,4-triazolyl, 1,3,4-triazolyl, pyranlyl, indolyl, pyrimidyl, thiazolyl, pyrazinyl, pyridazinyl, pyridyl, 4-pyridonyl, quinolyl and 10 1-isoquinolonyl.

A "carbocyclyl" is a saturated, partially saturated or unsaturated, mono or bicyclic carbon ring that contains 3-12 atoms; wherein a -CH₂- group can optionally be replaced by a -C(O)-. Preferably "carbocyclyl" is a monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 9 or 10 atoms. Suitable values for "carbocyclyl" include cyclopropyl, cyclobutyl, 15 1-oxocyclopentyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, phenyl, naphthyl, tetralinyl, indanyl or 1-oxoindanyl. Particularly "carbocyclyl" is cyclopropyl, cyclobutyl, 1-oxocyclopentyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, phenyl or 1-oxoindanyl.

An example of "C₁₋₁₀alkanoyloxy" and "C₁₋₆alkanoyloxy" is acetoxy. Examples of 20 "C₁₋₁₀alkoxycarbonyl" and "C₁₋₆alkoxycarbonyl" include methoxycarbonyl, ethoxycarbonyl, *n*- and *t*-butoxycarbonyl. Examples of "C₁₋₁₀alkoxy" and "C₁₋₆alkoxy" include methoxy, ethoxy and propoxy. Examples of "C₁₋₁₀alkanoylamino" and "C₁₋₆alkanoylamino" include formamido, acetamido and propionylamino. Examples of "C₁₋₆alkanoyl-*N*-(C₁₋₆alkyl)amino" include acetyl-*N*-methylamino and propionyl-*N*-ethyl-amino. Examples of "C₁₋₁₀alkylS(O)_a wherein a is 0 to 25 2" and "C₁₋₆alkylS(O)_a wherein a is 0 to 2" include methylthio, ethylthio, methylsulphinyl, ethylsulphinyl, mesyl and ethylsulphonyl. Examples of "C₁₋₁₀alkanoyl" and "C₁₋₆alkanoyl" include C₁₋₃alkanoyl, propionyl and acetyl. Examples of "*N*-(C₁₋₁₀alkyl)amino" and "*N*-(C₁₋₆alkyl)amino" include methylamino and ethylamino. Examples of "*N,N*-(C₁₋₁₀alkyl)₂amino" and "*N,N*-(C₁₋₆alkyl)₂amino" include di-*N*-methylamino, 30 di-(*N*-ethyl)amino and *N*-ethyl-*N*-methylamino. Examples of "C₂₋₁₀alkenyl" and "C₂₋₆alkenyl" are vinyl, allyl and 1-propenyl. Examples of "C₂₋₁₀alkynyl" and "C₂₋₆alkynyl" are ethynyl, 1-propynyl and 2-propynyl. Examples of "C₂₋₆alkylene" are ethylene, propylene and butylene.

- Examples of "C₂₋₆alkenyloxy" are vinyloxy, allyloxy and 1-propenyloxy. Examples of "N-(C₁₋₁₀alkyl)sulphamoyl" and "N-(C₁₋₆alkyl)sulphamoyl" are *N*-(C₁₋₃alkyl)sulphamoyl, *N*-(methyl)sulphamoyl and *N*-(ethyl)sulphamoyl. Examples of "N-(C₁₋₁₀alkyl)₂sulphamoyl" and "N-(C₁₋₆alkyl)₂sulphamoyl" are *N,N*-(dimethyl)sulphamoyl and
- 5 *N*-(methyl)-*N*-(ethyl)sulphamoyl. Examples of "N-(C₁₋₁₀alkyl)carbamoyl" and "N-(C₁₋₆alkyl)carbamoyl" are methylaminocarbonyl and ethylaminocarbonyl. Examples of "N,N-(C₁₋₁₀alkyl)₂carbamoyl" and "N,N-(C₁₋₆alkyl)₂carbamoyl" are dimethylaminocarbonyl and methylethylaminocarbonyl. Examples of "N-(C₁₋₁₀alkyl)carbamoyl" and "N-(C₁₋₆alkyl)carbamoyloxy" are methylaminocarbonyloxy and ethylaminocarbonyloxy.
- 10 Examples of "N,N-(C₁₋₁₀alkyl)₂carbamoyl" and "N,N-(C₁₋₆alkyl)₂carbamoyloxy" are dimethylaminocarbonyloxy and methylethylaminocarbonyloxy. Examples of "C₁₋₆alkylsulphonyl" are mesyl and ethylsulphonyl. Examples of "C₁₋₁₀alkylsulphonylamino" and "C₁₋₆alkylsulphonylamino" are mesylamino and ethylsulphonylamino. Examples of "C₁₋₆alkylsulphonyl-*N*-(C₁₋₆alkyl)amino" are mesyl-*N*-methylamino and ethylsulphonyl-*N*-
- 15 propylamino. Examples of "N'-(C₁₋₆alkyl)ureido" are *N'*-methylureido and *N'*-*i*-propylureido. Examples of "N-(C₁₋₆alkyl)ureido" are *N*-methylureido and *N*-*i*-propylureido. Examples of "N',N'-(C₁₋₆alkyl)₂ureido" are *N',N'*-dimethylureido and *N'*-methyl-*N'*-ethylureido. Examples of "N'-(C₁₋₆alkyl)-*N*-(C₁₋₆alkyl)ureido" are *N',N*-dimethylureido and *N'*-methyl-*N*-ethylureido. Examples of "N',N'-(C₁₋₆alkyl)₂-*N*-(C₁₋₆alkyl)ureido" are *N',N'*-dimethyl-*N*-methylureido and
- 20 *N'*-methyl-*N'*-ethyl-*N*-*t*-butylureido. Examples of "N,N,N-(C₁₋₁₀alkyl)₃ammonio" are trimethylamino and methyldiethylamino. Examples of "C₁₋₁₀alkoxycarbonylamino" and "C₁₋₆alkoxycarbonylamino" are methoxycarbonylamino and *t*-butoxycarbonylamino. Examples of "N-(C₁₋₁₀alkyl)sulphamoylamino" are *N*-methylsulphamoylamino and *N*-ethylsulphamoylamino. Examples of "N,N-(C₁₋₁₀alkyl)₂sulphamoylamino" are *N,N*-
- 25 dimethylsulphamoylamino and *N*-methyl-*N*-ethylsulphamoylamino. Examples of "carbocyclylC₁₋₁₀alkyl" include benzyl and phenethyl. Examples of "heterocyclylC₁₋₁₀alkyl" include 2-morpholinopropyl and pyridylmethyl. Examples of "phenylC₁₋₆alkoxy" include 2-phenylethoxy and 2-phenylpropoxy.

A suitable pharmaceutically acceptable salt of a compound of the invention is, for

30 example, an acid-addition salt of a compound of the invention which is sufficiently basic, for example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic, citric, acetate or maleic acid.

In addition a suitable pharmaceutically acceptable salt of a compound of the invention which is sufficiently acidic is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an organic base which affords a physiologically-acceptable cation, for example a salt with

5 methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

The compounds of the formula (I) may be administered in the form of a pro-drug which is broken down in the human or animal body to give a compound of the formula (I). examples of pro-drugs include *in vivo* hydrolysable esters and *in vivo* hydrolysable amides of a

10 compound of the formula (I).

An *in vivo* hydrolysable ester of a compound of the formula (I) containing carboxy or hydroxy group is, for example, a pharmaceutically acceptable ester which is hydrolysed in the human or animal body to produce the parent acid or alcohol. Suitable pharmaceutically acceptable esters for carboxy include C₁₋₆alkoxymethyl esters for example methoxymethyl,

15 C₁₋₆alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl esters, C₃₋₈cycloalkoxycarbonyloxyC₁₋₆alkyl esters for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters for example 5-methyl-1,3-dioxolen-2-onylmethyl; and C₁₋₆alkoxycarbonyloxyethyl esters for example 1-methoxycarbonyloxyethyl and may be formed at any carboxy group in the compounds of this invention.

20 An *in vivo* hydrolysable ester of a compound of the formula (I) containing a hydroxy group includes inorganic esters such as phosphate esters and α -acyloxyalkyl ethers and related compounds which as a result of the *in vivo* hydrolysis of the ester breakdown to give the parent hydroxy group. Examples of α -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxy-methoxy. A selection of *in vivo* hydrolysable ester forming groups

25 for hydroxy include alkanoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and N-(dialkylaminoethyl)-N-alkylcarbamoyl (to give carbamates), dialkylaminoacetyl and carboxyacetyl. Examples of substituents on benzoyl include morpholino and piperazino linked from a ring nitrogen atom via a methylene group to the 3- or 4- position of the benzoyl ring.

30 A suitable value for an *in vivo* hydrolysable amide of a compound of the formula (I) containing a carboxy group is, for example, a N-C₁₋₆alkyl or N,N-di-C₁₋₆alkyl amide such as N-methyl, N-ethyl, N-propyl, N,N-dimethyl, N-ethyl-N-methyl or N,N-diethyl amide.

Some compounds of the formula (I) may have chiral centres and/or geometric isomeric centres (E- and Z- isomers), and it is to be understood that the invention encompasses all such optical, diastereoisomers and geometric isomers that possess IBAT inhibitory activity.

The invention relates to any and all tautomeric forms of the compounds of the formula 5 (I) that possess IBAT inhibitory activity.

It is also to be understood that certain compounds of the formula (I) can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which possess IBAT inhibitory activity.

10 Particular values are as follows. Such values may be used where appropriate with any of the definitions, claims or embodiments defined hereinbefore or hereinafter.

M^1 is $-\text{CH}_2-$ and M^2 is $-\text{CR}^{22}\text{R}^{23}-$.

M^1 is $-\text{CH}_2-$ and M^2 is $-\text{NR}^{24}-$.

M^1 is $-\text{NR}^{21}-$ and M^2 is $-\text{CR}^{22}\text{R}^{23}-$.

15 R^{21} is hydrogen.

R^{21} is C_{1-6} alkyl.

R^{22} and R^{23} are independently selected from hydrogen and hydroxy.

One of R^{22} and R^{23} is hydrogen and the other is hydroxy.

R^{24} is hydrogen.

20 R^1 and R^2 are C_{1-4} alkyl.

R^1 and R^2 are both butyl.

One of R^1 and R^2 is ethyl and the other is butyl.

v is 0.

R^3 is C_{1-4} alkyl.

25 R^5 is a group of formula (IA).

R^6 is a group of formula (IA).

R^4 and R^7 are hydrogen.

R^5 is methylthio and R^6 is a group of formula (IA).

The R^5 or R^6 not selected from a group of formula (IA) is hydrogen, halo, C_{1-4} alkyl or 30 C_{1-4} alkylS(O)_a wherein a is 0.

The R^5 or R^6 not selected from a group of formula (IA) is hydrogen, bromo, methyl or methylthio.

The R⁵ or R⁶ not selected from a group of formula (IA) is hydrogen or methylthio.

The R⁵ or R⁶ not selected from a group of formula (IA) is hydrogen.

The R⁵ or R⁶ not selected from a group of formula (IA) is methylthio.

The group of formula (IA) is selected from:

- 5 *N*-{(R)- α -[N'-(2-hydroxyethyl)carbamoyl]benzyl}carbamoylmethoxy;
N-{(R)- α -[N'-(2-trimethylaminoethyl)carbamoyl]benzyl}carbamoylmethoxy;
N-{(R)- α -[N'-(2-aminoethyl)carbamoyl]benzyl}carbamoylmethoxy;
N-{(R)- α -[N'-(carbamoylmethyl)carbamoyl]benzyl}carbamoylmethoxy;
N-{(R)- α -[N'-((S)-1-carbamoyl-2-hydroxyethyl)carbamoyl]benzyl}carbamoylmethoxy;
- 10 *N*-((R)- α -carbamoylbenzyl)carbamoylmethoxy;
N-{(R)- α -[N'-(1,1-di-hydroxymethyl-2-hydroxyethyl)carbamoyl]benzyl}carbamoylmethoxy;
N-{(R)- α -[N'-(hydroxycarbamoylmethyl)carbamoyl]benzyl}carbamoylmethoxy;
N-((R)- α -{N'-[N-(2,2,2-trifluoroethyl)carbamoylmethyl]carbamoyl}benzyl)
carbamoylmethoxy;
- 15 *N*-((R)- α -{N'-[N-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoylmethyl]
carbamoyl}benzyl)carbamoylmethoxy;
N-((R)- α -{N'-[N-(2-fluoroethyl)carbamoylmethyl]carbamoyl}benzyl)carbamoylmethoxy;
N-((R)- α -{N'-[N-(ethyl)carbamoylmethyl]carbamoyl}benzyl)carbamoylmethoxy;
N-((R)- α -{N'-[N-(4-hydroxy-3-methoxybenzyl)carbamoylmethyl]carbamoyl}benzyl)
carbamoylmethoxy;
- 20 *N*-((R)- α -{N'-[N-(2-methoxyethyl)carbamoylmethyl]carbamoyl}benzyl)carbamoylmethoxy;
N-((R)- α -{N'-[N-(4-sulphamoylphenethyl)carbamoylmethyl]carbamoyl}benzyl)
carbamoylmethoxy;
N-((R)- α -{N'-[N-(2-*N,N*-dimethylaminosulphamoyl)ethyl]carbamoylmethyl]carbamoyl}
benzyl)carbamoylmethoxy;
- 25 *N*-[(R)- α -(N'-[N-(2-(N'-pyrimidin-2-ylureido)ethyl]carbamoylmethyl}carbamoyl)benzyl]
carbamoylmethoxy;
(*N*-{(R)- α -[N'-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]benzyl}carba
moylmethoxy;
- 30 *N*-{(R)- α -[N'-(3-morpholinopropyl)carbamoyl]benzyl}carbamoylmethoxy;
N-{(R)- α -[N'-(2-imidazol-4-ylethyl)carbamoyl]benzyl}carbamoylmethoxy;

- N*-{(R)- α -[*N'*-(2-*N,N*-dimethylaminosulphamoyl)ethyl]carbamoyl}benzyl} carbamoylmethoxy;
N-{(R)- α -[*N'*-(2-(2-hydroxyphenoxy)ethyl]carbamoyl}benzyl} carbamoylmethoxy;
N-{(R)- α -[*N'*-(3-hydroxy-1,5-benzodioxepin-3-ylmethyl)carbamoyl]benzyl}
 carbamoylmethoxy;
- 5 *N*-{(R)- α -[*N'*-(3-*t*-butoxycarbonylamino)benzyl]carbamoyl}benzyl} carbamoylmethoxy;
N-{(R)- α -[*N'*-(3-(benzyloxycarbonylimino-1-aminomethyl)benzyl]carbamoyl}benzyl}
 carbamoylmethoxy;
N-{(R)- α -[*N'*-(2-(3,4-dihydroxyphenyl)-2-methoxyethyl]carbamoyl}benzyl}
 carbamoylmethoxy;
- 10 *N*-{(R)- α -[*N'*-(2,3-dihydroxypropyl)carbamoyl]benzyl} carbamoylmethoxy;
N-{(R)- α -[*N'*-(2-(5-methoxyindol-3-yl)ethyl]carbamoyl}benzyl} carbamoylmethoxy;
N-{(R)- α -[*N'*-(2-(2,5-dioxothiazolidin-1-yl)ethyl]carbamoyl}benzyl} carbamoylmethoxy;
N-{(R)- α -[*N'*-(3-(4-methylpiperazin-1-yl)propyl]carbamoyl}benzyl} carbamoylmethoxy;
N-{(R)- α -[*N'*-(4-sulphamoylphenethyl)carbamoyl]benzyl} carbamoylmethoxy;
- 15 *N*-{(R)- α -[*N'*-(5,6-dimethoxy-2,3-dihydrobenzofuran-2-ylmethyl)carbamoyl]benzyl}
 carbamoylmethoxy;
N-{(R)- α -[*N'*-(1-*t*-butoxycarbonylpiperidin-4-ylmethyl)carbamoyl]benzyl} carbamoylmethoxy;
N-{(R)- α -[*N'*-(4-nitroanilino)carbamoylmethyl]carbamoyl}benzyl} carbamoylmethoxy;
N-{(R)- α -[*N'*-(2-(*N'*-pyrimidin-2-ylureido)ethyl]carbamoyl}benzyl} carbamoylmethoxy;
- 20 *N*-{(R)- α -[*N'*-(2-(*N'*-pyridin-2-ylureido)ethyl]carbamoyl}benzyl} carbamoylmethoxy;
N-{(R)- α -[*N'*-(2-(4-carbamoylphenoxy)ethyl]carbamoyl}benzyl} carbamoylmethoxy;
N-{(R)- α -[*N'*-(2-(2-oxoimidazolidin-1-yl)ethyl]carbamoyl}benzyl} carbamoylmethoxy; and
N-{(R)- α -[*N'*-(3-aminobenzyl)carbamoyl]benzyl} carbamoylmethoxy.

Z is -O- or -S(O)_b-; wherein b is 0.

25 R⁸ is hydrogen.

R⁹ is hydrogen.

R¹⁰ and R¹¹ are independently selected from hydrogen or carbocyclyl; wherein R¹⁰ and R¹¹ may be independently optionally substituted on carbon by one or more substituents selected from R²⁸.

30 R¹⁰ and R¹¹ are independently selected from hydrogen, cyclohexyl or phenyl; wherein R¹⁰ and R¹¹ may be independently optionally substituted on carbon by one or more substituents selected from halo.

R^{10} and R^{11} are independently selected from hydrogen, cyclohexyl or 2-fluorophenyl.

R^{13} is a group of formula (IB).

R^{14} is hydrogen.

R^{15} is hydrogen.

5 R^{17} is C_{1-10} alkyl; wherein R^{17} may be optionally substituted on carbon by one or more substituents selected from R^{47} or R^{17} is a group of formula (IC) (as depicted above).

R^{17} is C_{1-10} alkyl; wherein R^{17} may be optionally substituted on carbon by one or more substituents selected from R^{47} or R^{17} is a group of formula (IC) (as depicted above).

10 R^{17} is C_{1-6} alkyl; wherein R^{17} may be optionally substituted on carbon by one or more hydroxy or R^{17} is a group of formula (IC) (as depicted above).

R^{17} is pentyl substituted by 5 hydroxy or R^{17} is a group of formula (IC) (as depicted above).

R^{18} is selected from hydrogen.

R^{19} is selected from hydrogen.

15 R^{20} is C_{1-10} alkyl; wherein R^{20} may be independently optionally substituted on carbon by one or more R^{57} .

R^{20} is C_{1-10} alkyl; wherein R^{20} may be independently optionally substituted on carbon by one or more hydroxy.

R^{20} is pentyl substituted by 5 hydroxy.

20 p is 1.

q is 0.

r is 0.

m is 0.

n is 1.

25 z is 1.

R^{28} is halo.

R^{28} is fluoro.

R^{47} is hydroxy.

R^{57} is hydroxy.

30 One of R^5 and R^6 is a group of formula (IA) (as depicted above); wherein:

Z is -O- or -S(O)_b-; wherein b is 0;

R^8 is hydrogen;

R⁹ is hydrogen;

R¹⁰ and R¹¹ are independently selected from hydrogen or carbocyclyl; wherein R¹⁰ and R¹¹ may be independently optionally substituted on carbon by one or more substituents selected from R²⁸;

5 R¹³ is a group of formula (IB) (as depicted above);

R¹⁴ is hydrogen;

R¹⁵ is hydrogen;

R¹⁷ is C₁₋₁₀alkyl; wherein R¹⁷ may be optionally substituted on carbon by one or more substituents selected from R⁴⁷; or R¹⁷ is a group of formula (IC) (as depicted above) wherein:

10 R¹⁸ is selected from hydrogen;

R¹⁹ is selected from hydrogen;

R²⁰ is C₁₋₁₀alkyl; wherein R²⁰ may be independently optionally substituted on carbon by one or more R⁵⁷;

p is 1;

15 q is 0;

r is 0;

m is 0;

n is 1;

z is 1; and

20 R²⁸, R⁴⁷ and R⁵⁷ are independently selected from halo and hydroxy.

One of R⁵ and R⁶ is a group of formula (IA) (as depicted above); wherein:

Z is -O- or -S(O)_b-; wherein b is 0;

R⁸ is hydrogen;

R⁹ is hydrogen;

25 R¹⁰ and R¹¹ are independently selected from hydrogen, phenyl or carbocyclyl; wherein

R¹⁰ and R¹¹ may be independently optionally substituted on carbon by one or more substituents selected from halo;

R¹³ is a group of formula (IB) (as depicted above);

R¹⁴ is hydrogen;

30 R¹⁵ is hydrogen;

R^{17} is C_{1-6} alkyl; wherein R^{17} may be optionally substituted on carbon by one or more substituents selected from hydroxy; or R^{17} is a group of formula (IC) (as depicted above) wherein:

R^{18} is selected from hydrogen;

5 R^{19} is selected from hydrogen;

R^{20} is C_{1-6} alkyl; wherein R^{20} may be independently optionally substituted on carbon by one or more hydroxy;

p is 1;

q is 0;

10 r is 0;

m is 0;

n is 1; and

z is 1.

One of R^5 and R^6 is a group of formula (IA) (as depicted above); wherein:

15 Z is -O- or -S(O)_b-; wherein b is 0;

R^8 is hydrogen;

R^9 is hydrogen;

R^{10} and R^{11} are independently selected from hydrogen, 2-fluorophenyl or carbocyclyl;

R^{13} is a group of formula (IB) (as depicted above);

20 R^{14} is hydrogen;

R^{15} is hydrogen;

R^{17} is pentyl substituted by 5 hydroxy; or R^{17} is a group of formula (IC) (as depicted above) wherein:

R^{18} is selected from hydrogen;

25 R^{19} is selected from hydrogen;

R^{20} is pentyl substituted by 5 hydroxy;

p is 1;

q is 0;

r is 0;

30 m is 0;

n is 1; and

z is 1.

Therefore in an further aspect of the invention, there is provided a compound of formula (I) wherein:

M^1 is $-\text{CH}_2-$;

M^2 is $-\text{CR}^{22}\text{R}^{23}-$ and $-\text{NR}^{24}-$;

5 R^{22} and R^{23} are independently selected from hydrogen and hydroxy;

R^1 and R^2 are C_{1-4} alkyl;

v is 0;

R^4 and R^7 are hydrogen;

One of R^5 or R^6 is selected from a group of formula (IA) (as depicted above) and the

10 other is hydrogen or methylthio;

Z is $-\text{O}-$ or $-\text{S}(\text{O})_b-$; wherein b is 0;

R^8 is hydrogen;

R^9 is hydrogen;

R^{10} and R^{11} are independently selected from hydrogen or carbocyclyl; wherein R^{10} and

15 R^{11} may be independently optionally substituted on carbon by one or more substituents selected from R^{28} ;

R^{13} is a group of formula (IB) (as depicted above);

R^{14} is hydrogen;

R^{15} is hydrogen;

20 R^{17} is C_{1-10} alkyl; wherein R^{17} may be optionally substituted on carbon by one or more substituents selected from R^{47} ; or R^{17} is a group of formula (IC) (as depicted above) wherein:

R^{18} is selected from hydrogen;

R^{19} is selected from hydrogen;

R^{20} is C_{1-10} alkyl; wherein R^{20} may be independently optionally substituted on carbon by

25 one or more R^{57} ;

p is 1;

q is 0;

r is 0;

m is 0;

30 n is 1;

z is 1; and

R^{28} , R^{47} and R^{57} are independently selected from halo and hydroxy;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Therefore in an further aspect of the invention, there is provided a compound of formula (I) wherein:

M^1 is $-CH_2-$;

5 M^2 is $-CR^{22}R^{23}-$ and $-NR^{24}-$;

R^{22} and R^{23} are independently selected from hydrogen and hydroxy;

One of R^1 and R^2 is ethyl and the other is butyl;

v is 0;

R^4 and R^7 are hydrogen;

10 One of R^5 or R^6 is selected from a group of formula (IA) (as depicted above) and the other is hydrogen or methylthio;

Z is $-O-$ or $-S(O)_b-$; wherein b is 0;

R^8 is hydrogen;

R^9 is hydrogen;

15 R^{10} and R^{11} are independently selected from hydrogen, 2-fluorophenyl or carbocyclyl;

R^{13} is a group of formula (IB) (as depicted above);

R^{14} is hydrogen;

R^{15} is hydrogen;

20 R^{17} is pentyl substituted by 5 hydroxy; or R^{17} is a group of formula (IC) (as depicted above) wherein:

R^{18} is selected from hydrogen;

R^{19} is selected from hydrogen;

R^{20} is pentyl substituted by 5 hydroxy;

p is 1;

25 q is 0;

r is 0;

m is 0;

n is 1; and

z is 1;

30 or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.


In another aspect of the invention, preferred compounds of the invention are any one of the examples or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug

thereof.


Preferred aspects of the invention are those which relate to the compound of formula (I) or a pharmaceutically acceptable salt thereof.

Another aspect of the present invention provides a process for preparing a compound of formula (I) or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof which process (wherein variable groups are, unless otherwise specified, as defined in formula (I)) comprises of:

Process 1): for compounds of formula (I) wherein Z is -O-, -NR^a or -S-; reacting a compound of formula (IIa) or (IIb):




(IIa)



(IIb)


with a compound of formula (III):




(III)

wherein L is a displaceable group;

Process 2): reacting an acid of formula (IVa) or (IVb):




(IVa)




(IVb)

or an activated derivative thereof; with an amine of formula (V):




(V);

Process 3): for compounds of formula (I) wherein R^{13} is a group of formula (IB); reacting an acid of formula (VIa):




(VIa)

or (VIb):




(VIb)

with an amine of formula:




(VI)

Process 4): for compounds of formula (I) wherein R¹³ is a group of formula (IB) and R¹⁷ is a group of formula (IC); reacting an acid of formula (VIIIa):




(VIIIa)

or (VIIIb)




(VIIIb)

or an activated derivative thereof; with an amine of formula (IX):




(IX)

Process 5) for compounds of formula (I) wherein one of R⁵ and R⁶ are independently selected from C₁₋₆alkylthio optionally substituted on carbon by one or more R²⁵; reacting a compound of formula (Xa) or (Xb):



(Xa)



(Xb)

wherein L is a displaceable group; with a thiol of formula (XI):



(XI)


wherein R^m is C₁₋₆alkylthio optionally substituted on carbon by one or more R²⁵; and thereafter if necessary or desirable:

- i) converting a compound of the formula (I) into another compound of the formula (I);
- ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug.


L is a displaceable group, suitable values for L are for example, a halogeno or
5 sulphonyloxy group, for example a chloro, bromo, methanesulphonyloxy or
toluene-4-sulphonyloxy group.

Specific reaction conditions for the above reactions are as follows.


The bicyclic ring systems of the present invention may be assembled according to
Scheme Ia (1,4-benzothiazepine) or Scheme Ib (benzothiepinines) or Scheme IC (1,2-
10 benzothiazepines). The skilled person will appreciate to make any of the above identified
intermediates the value of R^4 or R^5 in the following schemes would be replaced with the
appropriate group. For example, to synthesize a compound of formula (IIa) R^4 would be HX
in the following schemes.



Scheme 1a



Scheme 1b



Scheme 1c

Starting materials for the above schemes are commercially available, or they are known in the literature, or they may be prepared by standard processes known in the art.

Process 1): Compounds of formula (IIa) or (IIb) may be reacted with compounds of formula (III) in the presence of a base for example an inorganic base such as sodium carbonate, or an organic base such as Hunigs base, in the presence of a suitable solvent such as acetonitrile, dichloromethane or tetrahydrofuran at a temperature in the range of 0°C to reflux, preferably at or near reflux.

Compounds of formula (III) are commercially available compounds, or they are known in the literature, or they are prepared by standard processes known in the art.

10 *Process 2*) and *Process 3*) and *Process 4*): Acids and amines may be coupled together in the presence of a suitable coupling reagent. Standard peptide coupling reagents known in the art can be employed as suitable coupling reagents, or for example carbonyldiimidazole and dicyclohexyl-carbodiimide, optionally in the presence of a catalyst such as dimethylaminopyridine or 4-pyrrolidinopyridine, optionally in the presence of a base for
15 example triethylamine, pyridine, or 2,6-di-*alkyl*-pyridines such as 2,6-lutidine or 2,6-di-*tert*-butylpyridine. Suitable solvents include dimethylacetamide, dichloromethane, benzene, tetrahydrofuran and dimethylformamide. The coupling reaction may conveniently be performed at a temperature in the range of -40 to 40°C.

Suitable activated acid derivatives include acid halides, for example acid chlorides, and
20 active esters, for example pentafluorophenyl esters. The reaction of these types of compounds with amines is well known in the art, for example they may be reacted in the presence of a base, such as those described above, and in a suitable solvent, such as those described above. The reaction may conveniently be performed at a temperature in the range of -40 to 40°C.

Amines of formula (V), (VII) and (IX) are commercially available, or they are known
25 in the literature, or they may be prepared by standard processes known in the art.

Process 4): Compounds of formula (Xa) and (Xb) may be reacted with thiols in the presence of base, for example an inorganic base such as sodium carbonate or an organic base such as Hunigs base, in the presence of a suitable solvent such as DMF or THF at a temperature in the range of 0°C to reflux.

30 Compounds of formula (Xa) and (Xb) may be prepared by any of the procedures above for the preparation of compounds of formula (I), but wherein one of R⁴ and R⁵ is L.

Compounds of formula (XI) are commercially available compounds, or they are known in the literature, or they are prepared by standard processes known in the art.

It will be appreciated that certain of the various ring substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or
5 generated by conventional functional group modifications either prior to or immediately following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such
10 procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions;
15 and the introduction of a halogeno group. Particular examples of modifications include the reduction of a nitro group to an amino group by for example, catalytic hydrogenation with a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating; oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl.

It will also be appreciated that in some of the reactions mentioned herein it may be
20 necessary/desirable to protect any sensitive groups in the compounds. The instances where protection is necessary or desirable and suitable methods for protection are known to those skilled in the art.

Conventional protecting groups may be used in accordance with standard practice (for illustration see T.W. Green, Protective Groups in Organic Synthesis, John Wiley and Sons,
25 1999). Thus, if reactants include groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or *t*-butoxycarbonyl group, an arylmethoxycarbonyl group,
30 for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an

an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

10 A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with
15 a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with
20 a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

The protecting groups may be removed at any convenient stage in the synthesis using
25 conventional techniques well known in the chemical art.

As stated hereinbefore the compounds defined in the present invention possess IBAT inhibitory activity. These properties may be assessed, for example, using an *in vitro* test assay for studying the effect on bile acid uptake in IBAT-transfected cells (Smith L., Price-Jones M. J., Hugnes K. T. and Jones N. R. A.; J Biomolecular Screening, 3, 227-230) or *in vivo* by
30 studying the effect on radiolabelled bile acid absorption in mice/rats (Lewis M. C., Brieady L. E. and Root C., J., J Lip Res 1995, 36, 1098-1105).

According to a further aspect of the invention there is provided a pharmaceutical

composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore in association with a pharmaceutically-acceptable diluent or carrier.

The composition may be in a form suitable for oral administration, for example as a
5 tablet or capsule, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) as a sterile solution, suspension or emulsion, for topical administration as an ointment or cream or for rectal administration as a suppository.

In general the above compositions may be prepared in a conventional manner using conventional excipients.

10 The compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, will normally be administered to a warm-blooded animal at a unit dose within the range 5-5000 mg per square meter body area of the animal, i.e. approximately 0.02-100 mg/kg, preferably 0.02 -50 mg/kg, and this normally provides a therapeutically-effective dose. A unit dose form such as a tablet or capsule will usually contain,
15 for example 1-250 mg of active ingredient. Preferably a daily dose in the range of 1-50 mg/kg, particularly 0.1-10 mg/kg is employed. In another aspect a daily dose in the range of 0.02-20 mg/kg is employed. However the daily dose will necessarily be varied depending upon the host treated, the particular route of administration, and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any
20 particular patient.

According to a further aspect of the present invention there is provided a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore for use in a method of prophylactic or therapeutic treatment of a warm-blooded animal, such as man.

25 We have found that the compounds defined in the present invention, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, are effective IBAT inhibitors, and accordingly have value in the treatment of disease states associated with hyperlipidaemic conditions.

Thus according to this aspect of the invention there is provided a compound of the
30 formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore for use as a medicament.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore in the manufacture of a medicament for use in the production of an IBAT inhibitory effect in a warm-blooded animal, such as man.

- 5 According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore in the manufacture of a medicament for use in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man.

- According to another feature of the invention there is provided the use of a compound
10 of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore in the manufacture of a medicament for use in the treatment of dyslipidemic conditions and disorders such as hyperlipidaemia, hypertriglyceridemia, hyperbetalipoproteinemia (high LDL), hyperprebetalipoproteinemia (high VLDL), hyperchylomicronemia, hypolipoproteinemia, hypercholesterolemia,
15 hyperlipoproteinemia and hypoalphalipoproteinemia (low HDL) in a warm-blooded animal, such as man.

- According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore in the manufacture of a medicament for use in the
20 treatment of different clinical conditions such as atherosclerosis, arteriosclerosis, arrhythmia, hyper-thrombotic conditions, vascular dysfunction, endothelial dysfunction, heart failure, coronary heart diseases, cardiovascular diseases, myocardial infarction, angina pectoris, peripheral vascular diseases, inflammation of cardiovascular tissues such as heart, valves, vasculature, arteries and veins, aneurisms, stenosis, restenosis, vascular plaques, vascular fatty
25 streaks, leukocytes, monocytes and/or macrophage infiltration, intimal thickening, medial thinning, infectious and surgical trauma and vascular thrombosis, stroke and transient ischaemic attacks in a warm-blooded animal, such as man.

- According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a
30 prodrug thereof, as defined hereinbefore in the manufacture of a medicament for use in the treatment of atherosclerosis, coronary heart diseases, myocardial infarction, angina pectoris,

peripheral vascular diseases, stroke and transient ischaemic attacks in a warm-blooded animal, such as man.

According to a further feature of this aspect of the invention there is provided a method for producing an IBAT inhibitory effect in a warm-blooded animal, such as man, in need of
5 such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to a further feature of this aspect of the invention there is provided a method of treating hyperlipidemic conditions in a warm-blooded animal, such as man, in need of such
10 treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to a further feature of this aspect of the invention there is provided a method of treating dyslipidemic conditions and disorders such as hyperlipidaemia, hypertriglyceridemia,
15 hyperbetalipoproteinemia (high LDL), hyperprebetalipoproteinemia (high VLDL), hyperchylomicronemia, hypolipoproteinemia, hypercholesterolemia, hyperlipoproteinemia and hypoalphalipoproteinemia (low HDL) in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug
20 thereof.

According to a further feature of this aspect of the invention there is provided a method of treating different clinical conditions such as atherosclerosis, arteriosclerosis, arrhythmia, hyper-thrombotic conditions, vascular dysfunction, endothelial dysfunction, heart failure, coronary heart diseases, cardiovascular diseases, myocardial infarction, angina pectoris,
25 peripheral vascular diseases, inflammation of cardiovascular tissues such as heart, valves, vasculature, arteries and veins, aneurisms, stenosis, restenosis, vascular plaques, vascular fatty streaks, leukocytes, monocytes and/or macrophage infiltration, intimal thickening, medial thinning, infectious and surgical trauma and vascular thrombosis, stroke and transient ischaemic attacks in need of such treatment which comprises administering to said animal an
30 effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to a further feature of this aspect of the invention there is provided a method of treating atherosclerosis, coronary heart diseases, myocardial infarction, angina pectoris, peripheral vascular diseases, stroke and transient ischaemic attacks in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

There is evidence that an IBAT inhibitor might potentially be useful in the treatment and/or prevention of gallstones. According to a further feature of this aspect of the invention there is provided a method of treating and / or preventing gallstones in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

The size of the dose required for the therapeutic or prophylactic treatment will necessarily be varied depending on the host treated, the route of administration and the severity of the illness being treated. A unit dose in the range, for example, 0.1-50mg/kg preferably 0.1-10 mg/kg is envisaged.

The IBAT inhibitory activity defined hereinbefore may be applied as a sole therapy or may involve, in addition to a compound of the invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. According to this aspect of the invention there is provided a pharmaceutical product comprising a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore and an additional IBAT inhibitory substance as defined hereinbefore and an additional hypolipidaemic agent for the conjoint treatment of hyperlipidaemia.

In another aspect of the invention, the compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, may be administered in association with an HMG Co-A reductase inhibitor, or pharmaceutically acceptable salts, solvates, solvates of such salts or prodrugs thereof. Suitable HMG Co-A reductase inhibitors, pharmaceutically acceptable salts, solvates, solvates of such salts or prodrugs thereof are statins well known in the art. Particular statins are fluvastatin, lovastatin, pravastatin, simvastatin, atorvastatin, cerivastatin, bervastatin, dalvastatin, mevastatin and

(E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulphonyl)amino]pyrimidin-5-yl](3R,5S)-3,5-dihydroxyhept-6-enoic acid (rosuvastatin), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof. A particular statin is atorvastatin, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof. A more particular statin is atorvastatin calcium salt. A further particular statin is (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulphonyl)amino]pyrimidin-5-yl](3R,5S)-3,5-dihydroxyhept-6-enoic acid (rosuvastatin), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof. A preferable particular statin is rosuvastatin calcium salt.

10 In an additional aspect of the invention, the compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof may be administered in association with an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and/or a bile acid binder thereby avoiding a possible risk of excess of bile acids in colon caused by the inhibition of the
15 ileal bile acid transport system. An excess of bile acids in the visceral contents may cause diarrhoea. Thus, the present invention also provides a treatment of a possible side effect such as diarrhoea in patients during therapy comprising the compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

An HMG CoA-reductase inhibitor, or a pharmaceutically acceptable salt, solvate,
20 solvate of such a salt or a prodrug thereof will by its action decrease the endogenous cholesterol available for the bile acid synthesis and have an additive effect in combination with the compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof on lipid lowering.

Suitable bile acid binders for such a combination therapy are resins, such as
25 cholestyramine and cholestipol. One advantage is that the dose of bile acid binder might be kept lower than the therapeutic dose for treatment of cholesterolaemia in single treatment comprising solely a bile acid binder. By a low dose of bile acid binder any possible side effects caused by poor tolerance of the patient to the therapeutic dose could also be avoided.

Therefore in an additional feature of the invention, there is provided a method for
30 producing an IBAT inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug

thereof in simultaneous, sequential or separate administration with an effective amount of an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Therefore in an additional feature of the invention, there is provided a method for
5 producing an IBAT inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with a bile acid binder.

Therefore in an additional feature of the invention, there is provided a method for
10 producing an IBAT inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such
15 a salt or a prodrug thereof, in simultaneous, sequential or separate administration with a bile acid binder.

Therefore in an additional feature of the invention, there is provided a method of treating hyperlipidemic conditions in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of
20 formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Therefore in an additional feature of the invention, there is provided a method of
25 treating hyperlipidemic conditions in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of a bile acid binder.

30 Therefore in an additional feature of the invention, there is provided a method of treating hyperlipidemic conditions in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of

formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in simultaneous, sequential or separate administration with a bile
5 acid binder.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug
10 thereof, in association with a pharmaceutically acceptable diluent or carrier.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a bile acid binder, in association with a pharmaceutically acceptable diluent or carrier.

15 According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a bile acid binder in association with a pharmaceutically acceptable diluent or
20 carrier.

According to a further aspect of the present invention there is provided a kit comprising a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

25 According to a further aspect of the present invention there is provided a kit comprising a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a bile acid binder.

According to a further aspect of the present invention there is provided a kit comprising a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate
30 of such a salt or a prodrug thereof, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof and a bile acid binder.

According to a further aspect of the present invention there is provided a kit comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a first unit dosage form;
- 5 b) an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof; in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

According to a further aspect of the present invention there is provided a kit comprising:

- 10 a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a first unit dosage form;
- b) a bile acid binder; in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

According to a further aspect of the present invention there is provided a kit

15 comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a first unit dosage form;
- b) an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof; in a second unit dosage form;
- 20 c) a bile acid binder; in a third unit dosage form; and
- d) container means for containing said first, second and third dosage forms.

According to a further aspect of the present invention there is provided a kit comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a
- 25 salt or a prodrug thereof, together with a pharmaceutically acceptable diluent or carrier, in a first unit dosage form;
- b) an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

30 According to a further aspect of the present invention there is provided a kit comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, together with a pharmaceutically acceptable diluent or carrier, in a first unit dosage form;
 - b) a bile acid binder, in a second unit dosage form; and
- 5 c) container means for containing said first and second dosage forms.

According to a further aspect of the present invention there is provided a kit comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, together with a pharmaceutically acceptable diluent or carrier, in a
- 10 first unit dosage form;
- b) an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a second unit dosage form; and
 - c) a bile acid binder; in a third unit dosage form; and
- d) container means for containing said first, second and third dosage forms.

15 According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the production of an IBAT inhibitory effect in a warm-blooded animal, such as man.

20 According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a bile acid binder, in the manufacture of a medicament for use in the production of an IBAT inhibitory effect in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound

25 of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a bile acid binder, in the manufacture of a medicament for use in the production of an IBAT inhibitory effect in a warm-blooded animal, such as man.

30 According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt,

solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a
5 prodrug thereof, a bile acid binder, in the manufacture of a medicament for use in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt,
10 solvate, solvate of such a salt or a prodrug thereof, and a bile acid binder, in the manufacture of a medicament for use in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man.

According to a further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula
15 (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the simultaneous, sequential or separate administration of an effective amount of an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier to a
20 warm-blooded animal, such as man in need of such therapeutic treatment.

According to a further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the
25 simultaneous, sequential or separate administration of an effective amount of a bile acid binder, optionally together with a pharmaceutically acceptable diluent or carrier to a warm-blooded animal, such as man in need of such therapeutic treatment.

According to a further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula
30 (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the simultaneous, sequential or separate administration of an effective amount of an HMG Co-A

reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable excipient, with the simultaneous, sequential or separate administration of an effective amount of a bile acid binder, optionally together with a pharmaceutically acceptable diluent or carrier to a warm-blooded
5 animal, such as man in need of such therapeutic treatment.

According to an additional further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with
10 the simultaneous, sequential or separate administration one or more of the following agents selected from:

- a CETP (cholesteryl ester transfer protein) inhibitor, for example those referenced and described in WO 00/38725 page 7 line 22 - page 10, line 17 which are incorporated herein by reference;
- 15 ➤ a cholesterol absorption antagonist for example azetidinones such as SCH 58235 and those described in US 5,767,115 which are incorporated herein by reference;
- a MTP (microsomal transfer protein) inhibitor for example those described in Science, 282, 751-54, 1998 which are incorporated herein by reference;
- a fibric acid derivative; for example clofibrate, gemfibrozil, fenofibrate, ciprofibrate and
20 bezafibrate;
- a nicotinic acid derivative, for example, nicotinic acid (niacin), acipimox and niceritrol;
- a phytosterol compound for example stanols;
- probucol;
- an anti-obesity compound for example orlistat (EP 129,748) and sibutramine (GB
25 2,184,122 and US 4,929,629);
- an antihypertensive compound for example an angiotensin converting enzyme inhibitor, an angiotensin II receptor antagonist, an adrenergic blocker, an alpha adrenergic blocker, a beta adrenergic blocker, a mixed alpha/beta adrenergic blocker, an adrenergic stimulant, calcium channel blocker, a diuretic or a vasodilator;
- 30 ➤ insulin;
- sulphonylureas including glibenclamide, tolbutamide;
- metformin; and/or

➤ acarbose;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier to a warm-blooded animal, such as man in need of such therapeutic treatment.

- 5 Particular ACE inhibitors or pharmaceutically acceptable salts, solvates, solvate of such salts or a prodrugs thereof, including active metabolites, which can be used in combination with a compound of formula (I) include but are not limited to, the following compounds: alacepril, alatriopril, altiopril calcium, ancovenin, benazepril, benazepril hydrochloride, benazeprilat, benzoylcaptopril, captopril, captopril-cysteine, captopril-glutathione, ceranapril, 10 ceranopril, ceronapril, cilazapril, cilazaprilat, delapril, delapril-diacid, enalapril, enalaprilat, enapril, epicaptopril, foroxymithine, fosfenopril, fosenopril, fosenopril sodium, fosinopril, fosinopril sodium, fosinoprilat, fosinoprilic acid, glycopril, hemorphan-4, idrapril, imidapril, indolapril, indolaprilat, libenzapril, lisinopril, lyciumin A, lyciumin B, mixanapril, moexipril, moexiprilat, moveltipril, muracein A, muracein B, muracein C, pentopril, perindopril, 15 perindoprilat, pivalopril, pivopril, quinapril, quinapril hydrochloride, quinaprilat, ramipril, ramiprilat, spirapril, spirapril hydrochloride, spiraprilat, spiropril, spiropril hydrochloride, temocapril, temocapril hydrochloride, teprotide,trandolapril,trandolaprilat, utibapril, zabicipril, zabiciprilat, zofenopril and zofenoprilat. Preferred ACE inhibitors for use in the present invention are ramipril, ramiprilat, lisinopril, enalapril and enalaprilat. More preferred 20 ACE inhibitors for uses in the present invention are ramipril and ramiprilat.

- Preferred angiotensin II antagonists, pharmaceutically acceptable salts, solvates, solvate of such salts or a prodrugs thereof for use in combination with a compound of formula (I) include, but are not limited to, compounds: candesartan, candesartan cilexetil, losartan, valsartan, irbesartan, tasosartan, telmisartan and eprosartan. Particularly preferred angiotensin 25 II antagonists or pharmaceutically acceptable derivatives thereof for use in the present invention are candesartan and candesartan cilexetil.

- In another aspect of the invention, the compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, may be administered in association with a PPAR alpha and/or gamma agonist, or pharmaceutically acceptable salts, 30 solvates, solvates of such salts or prodrugs thereof. Suitable PPAR alpha and/or gamma agonists, pharmaceutically acceptable salts, solvates, solvates of such salts or prodrugs thereof are well known in the art. These include the compounds described in WO 01/12187, WO

01/12612, WO 99/62870, WO 99/62872, WO 99/62871, WO 98/57941, WO 01/40170, J Med Chem, 1996, 39, 665, Expert Opinion on Therapeutic Patents, 10 (5), 623-634 (in particular the compounds described in the patent applications listed on page 634) and J Med Chem, 2000, 43, 527 which are all incorporated herein by reference. Particularly a PPAR alpha and/or
5 gamma agonist refers to WY-14643, clofibrate, fenofibrate, bezafibrate, GW 9578, troglitazone, pioglitazone, rosiglitazone, eglitazone, proglitazone, BRL-49634, KRP-297, JTT-501, SB 213068, GW 1929, GW 7845, GW 0207, L-796449, L-165041 and GW 2433. Particularly a PPAR alpha and/or gamma agonist refers to (S)-2-ethoxy-3-[4-(2-{4-methanesulphonyloxyphenyl}ethoxy)phenyl] propanoic acid and pharmaceutically acceptable
10 salts thereof. Additional suitable PPAR alpha and/or gamma agonists are NN622/Ragaglitazar and BMS 298585.

Therefore in an additional feature of the invention, there is provided a method for producing an IBAT inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of
15 formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Therefore in an additional feature of the invention, there is provided a method of
20 treating hyperlipidemic conditions in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of
25 such a salt or a prodrug thereof.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug
30 thereof, in association with a pharmaceutically acceptable diluent or carrier.

According to a further aspect of the present invention there is provided a kit comprising a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate

of such a salt or a prodrug thereof, and a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to a further aspect of the present invention there is provided a kit comprising:

- 5 a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a first unit dosage form;
- b) a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof; in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

10 According to a further aspect of the present invention there is provided a kit comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, together with a pharmaceutically acceptable diluent or carrier, in a first unit dosage form;
- 15 b) a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a
20 prodrug thereof, and a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the production of an IBAT inhibitory effect in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a
25 prodrug thereof, a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man.

According to a further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula
30 (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the simultaneous, sequential or separate administration of an effective amount of a PPAR alpha

and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier to a warm-blooded animal, such as man in need of such therapeutic treatment.

In addition to their use in therapeutic medicine, the compounds of formula (I), or a
5 pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effects of inhibitors of IBAT in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

Many of the intermediates described herein are novel and are thus provided as a further
10 feature of the invention. For example compounds of formula (IVa), (IVb), (VIa), (VIb), (VIIIa) and (VIIIb) show IBAT inhibitory activity when tested in the above referenced *in vitro* test assay.

Thus in a further feature of the invention, there is provided a compound of formula (IVa), (IVb), (VIa), (VIb), (VIIIa) or (VIIIb), or a pharmaceutically acceptable salt, solvate,
15 solvate of such a salt or a prodrug thereof; with the proviso that when R¹¹ and R¹² are selected from carbocyclyl and / or heterocyclyl, said carbocyclyl and heterocyclyl are not totally unsaturated.

Therefore according to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (IVa), (IVb), (VIa),
20 (VIb), (VIIIa) or (VIIIb), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore in association with a pharmaceutically-acceptable diluent or carrier; with the proviso that when R¹¹ and R¹² are selected from carbocyclyl and / or heterocyclyl, said carbocyclyl and heterocyclyl are not totally unsaturated.

According to an additional aspect of the present invention there is provided a
25 compound of the formula (IVa), (IVb), (VIa), (VIb), (VIIIa) or (VIIIb), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore for use in a method of prophylactic or therapeutic treatment of a warm-blooded animal, such as man; with the proviso that when R¹¹ and R¹² are selected from carbocyclyl and / or heterocyclyl, said carbocyclyl and heterocyclyl are not totally unsaturated.

30 Thus according to this aspect of the invention there is provided a compound of the formula (IVa), (IVb), (VIa), (VIb), (VIIIa) or (VIIIb), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore for use as a

medicament; with the proviso that when R^{11} and R^{12} are selected from carbocyclyl and / or heterocyclyl, said carbocyclyl and heterocyclyl are not totally unsaturated.

According to another feature of the invention there is provided the use of a compound of the formula (IVa), (IVb), (VIa), (VIb), (VIIIa) or (VIIIb), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof as defined hereinbefore in the manufacture of a medicament for use in the production of an IBAT inhibitory effect in a warm-blooded animal, such as man; with the proviso that when R^{11} and R^{12} are selected from carbocyclyl and / or heterocyclyl, said carbocyclyl and heterocyclyl are not totally unsaturated.

According to another feature of the invention there is provided the use of a compound of the formula (IVa), (IVb), (VIa), (VIb), (VIIIa) or (VIIIb), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof as defined hereinbefore in the manufacture of a medicament for use in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man; with the proviso that when R^{11} and R^{12} are selected from carbocyclyl and / or heterocyclyl, said carbocyclyl and heterocyclyl are not totally unsaturated.

According to a further feature of this aspect of the invention there is provided a method for producing an IBAT inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (IVa), (IVb), (VIa), (VIb), (VIIIa) or (VIIIb), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof; with the proviso that when R^{11} and R^{12} are selected from carbocyclyl and / or heterocyclyl, said carbocyclyl and heterocyclyl are not totally unsaturated.

According to a further feature of this aspect of the invention there is provided a method of treating hyperlipidemic conditions in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (IVa), (IVb), (VIa), (VIb), (VIIIa) or (VIIIb), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof; with the proviso that when R^{11} and R^{12} are selected from carbocyclyl and / or heterocyclyl, said carbocyclyl and heterocyclyl are not totally unsaturated.

In the above other pharmaceutical composition, process, method, use and medicament manufacture features, the alternative and preferred embodiments of the compounds of the invention described herein also apply.

Examples

The invention will now be illustrated in the following non limiting examples, in which standard techniques known to the skilled chemist and techniques analogous to those described in these examples may be used where appropriate, and in which, unless otherwise stated:

- 5 (i) evaporations were carried out by rotary evaporation in vacuo and work up procedures were carried out after removal of residual solids such as drying agents by filtration;
- (ii) all reactions were carried out under an inert atmosphere at ambient temperature, typically in the range 18-25°C, with solvents of HPLC grade under anhydrous conditions, unless otherwise stated;
- 10 (iii) column chromatography (by the flash procedure) was performed on Silica gel 40-63 μm (Merck);
- (iv) yields are given for illustration only and are not necessarily the maximum attainable;
- (v) the structures of the end products of the formula (I) were generally confirmed by nuclear (generally proton) magnetic resonance (NMR) and mass spectral techniques; magnetic
- 15 resonance chemical shift values were measured in deuterated CD_3OD (unless otherwise stated) on the delta scale (ppm downfield from tetramethylsilane); proton data is quoted unless otherwise stated; spectra were recorded on a Varian Mercury-300 MHz, Varian Unity plus-400 MHz, Varian Unity plus-600 MHz or on Varian Inova-500 MHz spectrometer; and peak multiplicities are shown as follows: s, singlet; d, doublet; dd, double doublet; t, triplet; tt,
- 20 triple triplet; q, quartet; tq, triple quartet; m, multiplet; br, broad; LCMS were recorded on a Waters ZMD, LC column xTerra MS C_8 (Waters), detection with a HP 1100 MS-detector diode array equipped; mass spectra (MS) (loop) were recorded on VG Platform II (Fisons Instruments) with a HP-1100 MS-detector diode array equipped; unless otherwise stated the mass ion quoted is (MH^+);
- 25 (vi) unless further details are specified in the text, analytical high performance liquid chromatography (HPLC) was performed on Prep LC 2000 (Waters), Kromasil C_8 , 7 μm , (Akzo Nobel); MeCN and de-ionised water 100 mM ammonium acetate as mobile phases, with suitable composition;
- (vii) intermediates were not generally fully characterised and purity was assessed by thin layer
- 30 chromatography (TLC), HPLC, infra-red (IR), MS or NMR analysis;
- (viii) where solutions were dried sodium sulphate was the drying agent;

(ix) where an "ISOLUTE" column is referred to, this means a column containing 2g of silica, the silica being contained in a 6ml disposable syringe and supported by a porous disc of 54Å pore size, obtained from International Sorbent Technology under the name "ISOLUTE";

"ISOLUTE" is a registered trade mark;

5 (x) the following abbreviations may be used hereinbefore or hereinafter:-

DCM	dichloromethane;
DMF	N,N-dimethylformamide;
TFA	trifluoroacetic acid;
TBTU	o-Benzotriazol-1-yl-N,N,N',N'-tetramethyluronium tetrafluoroborate;
10 EtOAc	ethyl acetate; and
MeCN	acetonitrile.

Example 1

15 (+/-)-trans-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(N-{(R)-α-[N-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine

(+/-)-trans-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-8-(carboxymethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine (Method 2; 20 mg, 0.043 mmol), (R)-α-[N-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]benzylamine (Method 10; 19 mg, 0.064 mmol) and N-methylmorpholine (27 mg, 0.26 mmol) were dissolved in DMF and stirred at room temperature. TBTU (19 mg, 0.059 mmol) was added and the reaction mixture was stirred for 2 hours. LC-MS indicated no unreacted starting material. A small amount of water and acetonitrile was added and the product was purified by preparative HPLC on a C8 column using a gradient from 5% to 100% MeCN in 0.1M ammonium acetate buffer as mobile phase.

25 Lyophilisation gave 27 mg of the title compound. NMR (400 MHz, CD₃OD): 0.81 (t, 3H), 0.89 (t, 3H), 1.11-1.35 (m, 4H), 1.4-1.5 (m, 1H), 1.52-1.63 (m, 1H), 1.74-1.85 (m, 1H), 1.94 (s, 3H), 2.27-2.28 (m, 1H), 3.11-3.25 (m, 2H), 3.46-3.53 (m, 2H), 3.53-3.75 (m, 5H), 3.78-3.84 (m, 1H), 4.63-4.66 (m, 1H), 5.51 (s, 1H), 6.0 (s, 1H), 6.59-6.64 (m, 1H), 6.95-7.01 (m, 1H), 7.27-7.35 (m, 4H), 7.35-7.41 (m, 6H), 7.66 (t, 1H).

Example 2

(+/-)-trans-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(N-{(R)- α -[N-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine

5 3,5-*trans*-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(N-{(R)- α -[N-(carboxymethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine (Method 6; 2mg, 3 μ mol) was dissolved in 0.5ml DMF. (D)-Glucamine (2mg, 11 μ mol), N-methyl morpholine (ca 1 μ l, 9 μ mol) and TBTU (1.1mg, 4 μ mol) were added and the solution was stirred for 3hours. The reaction was completed according to LC/MS-analysis. The product
10 was purified by preparative HPLC on a C8 column using a gradient from 5% to 100% MeCN in 0.1M ammonium acetate buffer as mobile phase. The product fraction was lyophilised to yield 2.4mg (95%). LC/Micromass Q TOF micro MS analysis showed four peaks corresponding to four isomers of the product with a total purity of 96%; m/z=831.328.

15 **Example 3**

1,1-Dioxo-3-ethyl-3-butyl-4-hydroxy-5-phenyl-7-(N-{ α -[N-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]-2-fluorobenzyl}carbamoylmethylthio)-2,3,4,5-tetrahydrobenzothiapiene

1,1-Dioxo-3-butyl-3-ethyl-4-hydroxy-5-phenyl-7-{N-[α -(carboxy)-2-fluorobenzyl]
20 carbamoylmethylthio}-2,3,4,5-tetrahydrobenzothiepine (Method 3; 28 mg, 0.046 mmol) and D-glucamine (10 mg, 0.055 mmol) were added to DMF (2.0 ml) and 2,6-lutidine (0.02 ml, 0.14 mmol) was added. When everything had dissolved TBTU (20 mg, 0.060 mmol) was added and the reaction mixture was stirred at ambient temperature overnight. More D-glucamine (10 mg, 0.055 mmol) and TBTU (20 mg, 0.060 mmol) were added and the reaction
25 mixture was stirred for another 3 hours. The reaction mixture was purified with preparative HPLC using MeCN/ammonium acetate buffer (gradient 50:50 to 100:0) to give the title compound (9.0 mg, 25 %). NMR (400 MHz, DMSO-d₆): 0.73 (t, 3 H), 0.83 (t, 3 H), 1.00 (m, 1 H), 1.20-1.45 (m, 5 H), 1.59 (m, 1 H), 2.01 (m, 1 H), 3.10 (m, 2 H), 3.94 (s, 1 H), 5.22 (d, 1 H), 5.60 (d, 1 H), 6.41 (dd, 1 H), 7.13 (m, 3 H), 7.25-7.43 (m, 8 H), 7.74 (dd, 1 H), 8.08 (br
30 s, 1 H), 8.89 (br s, 1 H).

Example 4

1,1-Dioxo-3-butyl-3-ethyl-4-hydroxy-5-phenyl-7-(N-{1-[N-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]-1-(cyclohexyl)methyl}carbamoylmethylthio)-2,3,4,5-tetrahydrobenzothiepine

5 1,1-Dioxo-3-butyl-3-ethyl-4-hydroxy-5-phenyl-7-{N-[1-(carboxy)-1-(cyclohexyl)methyl] carbamoylmethylthio }-2,3,4,5-tetrahydrobenzothiepine (Method 5; 4 mg, 0.0066 mmol), D-glucamine (2 mg, 0.0088 mmol) and N-methylmorpholine (4 mg, 0.035 mmol) were dissolved in DMF and stirred at room temperature for 10 min. TBTU (0.003 mg, 0.0087 mmol) and the reaction mixture was stirred over night. A small amount of water and
10 acetonitrile was added and the product was purified by preparative HPLC on a C8 column using a gradient from 5% to 100% MeCN in 0.1 %M ammonium acetate buffer as mobile phase. Lyophilization gave (1.7 mg 33.4%) of the title compound. LC/MicromassQ TOF micro MS analysis showed two peaks corresponding to two isomers of the product with a total purity of 96%; m/z=765.343

15

Preparation of Starting Materials

The starting materials for the Examples above are either commercially available or are readily prepared by standard Methods from known materials. For example, the following reactions are an illustration, but not a limitation, of some of the starting materials used in the
20 above reactions.

Method 1

(+/-)-trans-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-8-(t-butoxycarbonylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine

25 A mixture of (+/-)-trans-1,1-dioxo-3-butyl-3-ethyl-5-phenyl-8-hydroxy-2,3,4,5-tetrahydro-1,4-benzothiazepine (prepared according to WO/9605188; 52 mg, 0.14 mmol), t-butyl bromoacetate (32 mg, 0.16 mmol), potassium carbonate (76 mg, 0.55 mmol) and cesium carbonate (12 mg, 0.036 mmol) in acetonitrile was stirred at 50 C for 2 hours. The solvent was removed at reduced pressure and the crude product was purified by chromatography. The
30 eluent was initially DCM and thereafter switched to EtOAc:DCM, 1:10. 0.055 g of the desired product was obtained.

Method 2

(+/-)-trans-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-8-(carboxymethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine

(+/-)-trans-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-8-(*t*-butoxycarbonylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine (Method 1; 58 mg, 0.12 mmol) was dissolved in formic acid and stirred for 20 hours at ambient temperature and then for 4 hours at 45°C. The formic acid was removed at reduced pressure. Toluene was added to the residue and the toluene and remaining formic acid were removed at reduced pressure to give the desired product (50 mg, 0.12 mmol).

10 Method 3

1,1-Dioxo-3-butyl-3-ethyl-4-hydroxy-5-phenyl-7-{*N*-[α -(carboxy)-2-fluorobenzyl] carbamoylmethylthio}-2,3,4,5-tetrahydrobenzothiepine

1,1-Dioxo-3-butyl-3-ethyl-4-hydroxy-5-phenyl-7-(carboxymethylthio)-2,3,4,5-tetrahydrobenzothiepine (Method 4; 59 mg, 0.13 mmol) and methyl amino(2-fluorophenyl)acetate hydrochloride (40 mg, 0.18 mmol) were added to DCM (3.0 ml) and 2,6-lutidine (0.04 ml, 0.38 mmol) was added. The reaction mixture was stirred at ambient temperature for 5 min and TBTU (58 mg, 0.18 mmol) was added. After stirring at ambient temperature for 3 hours the reaction mixture was diluted with DCM (5 ml) and was washed with HCl (aq., 1 M, 5 ml). The solvent was evaporated and the residue was dissolved in THF (2.0 ml) and water (1.0 ml) and NaOH (aq., 1 M, 0.30 mmol) was added. After stirring at ambient temperature for 2 h the reaction mixture was quenched with HCl (aq., 1 M), diluted with water (10 ml) and extracted with DCM (3 x 5 ml). The combined organic layers were purified with preparative HPLC MeCN/ ammonium acetate buffer 40:60 to give the title compound (70 mg, 89 %). NMR (400 MHz, DMSO- d_6): 0.74 (t, 3 H), 0.84 (t, 3 H), 1.00 (m, 1 H), 1.18-1.46 (m, 5 H), 1.60 (m, 1 H), 2.02 (m, 1 H), 3.11 (m, 2 H), 3.69 (m, 2 H), 3.94 (s, 1 H), 5.04 (t, 1 H), 5.23 (d, 1 H), 6.44 (d, 1 H), 6.94-7.44 (m, 11 H), 7.70 (t, 1 H), 8.40 (m, 1 H).

Method 4**1,1-Dioxo-3-butyl-3-ethyl-4-hydroxy-5-phenyl-7-(carboxymethylthio)-2,3,4,5-tetrahydrobenzothiepine**

1,1-Dioxo-3-butyl-3-ethyl-4-hydroxy-5-phenyl-7-fluoro-2,3,4,5-

- 5 tetrahydrobenzothiepine (prepared according to WO 98/40375; 300 mg, 0.77 mmol) was dissolved in DMF (3 ml) and caesium carbonate (540 mg, 1.66 mmol) was added followed by ethyl thioglycolate (0.17 ml, 1.54 mmol) and the reaction mixture was subjected to microwave irradiation in a Smith Synthesiser at 100°C for 5 min. The reaction mixture was added to water (25 ml), extracted with DCM (5 x 5 ml) and concentrated. The crude ethyl ester was dissolved
- 10 in THF (4 ml) and water (2 ml) and NaOH (1 M, 4 ml) was added. The reaction mixture was stirred at ambient temperature for 6 hours, quenched with HCl (1 M), diluted with water (25 ml) and extracted with DCM (4 x 5 ml). The combined organic layers were purified with preparative HPLC using MeCN/ammonium acetate buffer as the eluent (50:50) to give the title compound (172 mg, 48 %). NMR (400 MHz, DMSO-d₆): 0.74 (t, 3H), 0.84 (t, 3H), 1.01 (m,
- 15 1H), 1.18-1.46 (m, 5H), 1.60 (m, 1H), 2.02 (m, 1H), 3.07/3.12 (ABq, 2H), 3.33/3.39 (ABq, 2H), 3.95 (s, 1H), 5.23 (s, 1H), 6.38 (d, 1H), 7.19 (dd, 1H), 7.34 (m, 1H), 7.42 (m, 4H), 7.73 (d, 1H).

Method 5

- 20 **1,1-Dioxo-3-butyl-3-ethyl-4-hydroxy-5-phenyl-7-{N-[1-(carboxy)-1-(cyclohexyl)methyl]carbamoyl}methylthio}-2,3,4,5-tetrahydrobenzothiepine**

The title compound was synthesised according to the same procedure as Method 3 but using methyl amino(cyclohexyl)acetate hydrochloride as starting material. NMR (400 MHz, DMSO-d₆): 0.74 (t, 3 H), 0.84 (t, 3 H), 0.92-1.10 (m, 6 H), 1.20-1.66 (m, 12 H), 2.01 (m, 1

25 H), 3.07/3.13 (ABq, 2 H), 3.60-3.75 (m, 3 H), 3.95 (m, 2 H), 5.24 (s, 1 H), 6.44 (m, 1 H), 7.33 (m, 1 H), 7.41 (m, 4 H), 7.75 (d, 1 H), 7.98 (t, 1 H).

Method 6

- 30 **3,5-trans-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(N-{(R)- α -[N-(carboxymethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine**

3,5-trans-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(carboxymethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine (Method 7; 50 mg, 0.105 mmol) was dissolved in

DCM (2 ml). 2,6-Lutidine (0.025 ml, 0.215 mmol), TBTU (45 mg, 0.140 mmol) and (R)- α -[N-(*t*-butoxycarbonylmethyl)carbamoyl]benzylamine (Method 4; 43 mg, 0.163 mmol) were added successively. The mixture was stirred for 2 hours at ambient temperature. The solution was concentrated and the intermediate ester was purified by chromatography on silica using 5 DCM/EtOAc (9/1) as eluent. The solvent was evaporated to yield 45 mg (60%). *M/z*: 724. The ester was dissolved in 3 ml DCM and hydrolysed by addition of TFA (1 ml). After 2 hours the mixture was concentrated and purified using preparative HPLC. A gradient of MeCN from 40% to 60% in 0.1 M ammonium acetate buffer was used as eluent. Lyophilisation yielded 33 mg (80%). NMR (400 MHz): 0.75-0.85 (m, 3H), 0.85-0.95 (m, 3H), 1.1-1.65 (m, 6H), 1.75-10 1.9 (m, 1H), 2.0 (s, 3H), 2.2-2.4 (m, 1H), 3.1-3.55 (m, 2H), 3.85 (ABq, 2H), 4.6-4.8 (m, 2H), 5.6 (s, 1H), 5.98-6.03 (m, 1H), 6.4 (s, 1H), 7.25-7.56 (m, 11H); *m/z*: 668.

Method 7

3,5-*trans*-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(carboxymethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine

The title compound was prepared from 3,5-*trans*-1,1-dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-hydroxy-2,3,4,5-tetrahydro-1,4-benzothiazepine (Method 8; 153 mg, 0.36 mmol), ethylbromoacetate (75 mg, 0.45 mmol), sodium carbonate (162 mg, 1.52 mmol) and tetrabutylammonium bromide (12 mg, 0.037 mmol) in acetonitrile (10 ml). The intermediate 20 ethyl ester was extracted between diluted HCl and DCM. The DCM phase was washed with brine, dried with Na₂SO₄ and concentrated. *M/z* 506. The crude product was dissolved in THF/H₂O (3/1; 4 ml) and LiOH (22 mg, 0.91 mmol) was added. The mixture was stirred for 2h and the solvent was removed under reduced pressure. The crude product was purified using 25 preparative HPLC. A gradient from 40% to 60% MeCN in 0.1 M ammonium acetate buffer was used as eluent. The MeCN was removed under reduced pressure and the remaining aqueous solution was acidified using 5% HCl and was then extracted with DCM. The DCM layer was dried with Na₂SO₄ and concentrated. The crude product was co-evaporated with diethyl ether. The obtained crystals were filtered off and dried. Mass: 158 mg (91%). NMR 0.75 (t, 3H), 0.9 (t, 3H), 1.1-1.7 (m, 6H), 1.7-1.9 (m, 1H), 2.0 (s, 3H), 2.2-2.4 (m, 1H), 3.3 30 (dd, 2H), 4.75 (s, 2H), 6.0 (s, 1H), 6.4 (s, 1H), 7.3-7.5 (m, 6H); *m/z*: 478.

Method 8**3,5-trans-1,1-Dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-hydroxy-2,3,4,5-tetrahydro-1,4-benzothiazepine**

(+/-)-trans-7-Bromo-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-

- 5 benzothiazepine 1,1-dioxide (prepared according to WO 96/05188; 300 mg, 0.64 mmol) was dissolved in 5 ml DMF under N₂(g)-atmosphere. Sodium thiomethylate (150 mg, 2.14 mmol) was added and the mixture was heated to 110°C for 2h. The solvent was removed under reduced pressure and the residue was extracted between 5% HCl and EtOAc. The organic phase was washed with brine, dried with Na₂SO₄ and concentrated. The product was purified
- 10 using preparative HPLC. A gradient from 40% to 100% of MeCN in 0.1 M ammonium acetate buffer was used as eluent. Lyophilisation yielded 153 mg, 57%. M/z: 420.

Method 9**(N-{(R)-α-[N-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-Pentahydroxyhexyl)carbamoyl]benzyl}**

- 15 carbamoyl)oxymethylphenyl

(2R)--[[(Benzyloxy)carbonyl]amino }(phenyl)acetic acid (2.07g, 7.26 mmol),

(2R,3R,4R,5S)-6-aminoheptane-1,2,3,4,5-pentol (1.44g, 7.9 mmol) and 2,6-dimethylpyridine (0.933g, 8.7 mmol) were dissolved in DMF (30 ml). TBTU (3.80g, 8.7 mmol) was added and the mixture stirred for 1.5 hour. (2R,3R,4R,5S)-6-aminoheptane-1,2,3,4,5-pentol (0.21g, 0.2

- 20 mmol) was added and the mixture stirred for 1 hour. The solvent was evaporated under reduced pressure. The mixture was dissolved in isopropanol (35 ml), heated to 80-100°C. The mixture was then allowed to reach room temperature. The formed crystals was filtered off and washed with isopropanol (2 x 20 ml) and two times with diethyl ether (2 x 25 ml). The crystals was recrystallized using isopropanol (100 ml) heated to reflux. The mixture was then allowed
- 25 to reach room temperature. The crystals was filtered off and washed with small portions of isopropanol (totally 170 ml) and with diethyl ether (2 x 50 ml) to yield 2.517g(77%) of the product. NMR (600 MHz, CD₃OD): 3.15-3.23 (m, 1H), 3.48 (dd, 1H), 3.53-3.68 (m, 4H), 3.71 (dd, 1H), 3.77-3.83 (m, 1H), 5.08 (s, 2H), 5.20 (s, 1H), 7.24-7.36 (m, 8H), 7.39 (brd, 2H)


Method 10

(R)- α -[N'-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-Pentahydroxyhexyl)carbamoyl]benzylamine

(N-{(R)- α -[N'-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-Pentahydroxyhexyl)carbamoyl]benzyl}carbamoyl)oxymethylphenyl (Method 9; 2.46g, 5.49 mmol), palladium on activated
5 carbon (5%, 0.535g), water (1.5 ml) was dissolved in ethanol (80 ml). The mixture was treated under H₂-atmosphere overnight. The mixture was filtered through hyflo and the solvent was evaporated under reduced pressure to yield 1.543g(89.5%) of the title compound.
NMR (400 MHz, CD₃OD): δ 3.23 (dd, 1H), 3.49 (dd, 1H), 3.56-3.62 (m, 2H), 3.63-3.70 (m, 2H), 3.73 (dd, 1H), 3.79-3.84 (m, 1H), 4.48 (s, 1H), 7.25-7.37 (m, 3H), 7.41 (brd, 2H).

Claim

1. A compound of formula (I):



(I)

5

wherein

M^1 is $-CH_2-$ or $-NR^{21}-$;

M^2 is $-CR^{22}R^{23}-$ or $-NR^{24}-$; provided that if M^1 is $-NR^{21}-$, M^2 is $-CR^{22}R^{23}-$;

One of R^1 and R^2 are selected from hydrogen, C_{1-6} alkyl or C_{2-6} alkenyl and the other is

10 selected from C_{1-6} alkyl or C_{2-6} alkenyl;

R^3 is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy,


$N-(C_{1-6}alkyl)amino$, $N,N-(C_{1-6}alkyl)_2amino$, $C_{1-6}alkanoylamino$, $N-(C_{1-6}alkyl)carbamoyl$,

$N,N-(C_{1-6}alkyl)_2carbamoyl$, $C_{1-6}alkylS(O)_a$ wherein a is 0 to 2, $C_{1-6}alkoxycarbonyl$,

15 $N-(C_{1-6}alkyl)sulphamoyl$ and $N,N-(C_{1-6}alkyl)_2sulphamoyl$;

v is 0-5;

one of R^5 and R^6 is a group of formula (IA):



(IA)

20

R^4 and R^7 and the other of R^5 and R^6 are independently selected from hydrogen, halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, C_{1-4} alkanoyl, C_{1-4} alkanoyloxy, $N-(C_{1-4}alkyl)amino$, $N,N-(C_{1-4}alkyl)_2amino$, $C_{1-4}alkanoylamino$, $N-(C_{1-4}alkyl)carbamoyl$, $N,N-(C_{1-4}alkyl)_2carbamoyl$,

$C_{1-4}alkylS(O)_a$ wherein a is 0 to 2, $C_{1-4}alkoxycarbonyl$, $N-(C_{1-4}alkyl)sulphamoyl$ and $N,N-(C_{1-4}alkyl)_2sulphamoyl$; wherein R^4 and R^7 and the other of R^5 and R^6 may be optionally substituted on carbon by one or more R^{25} ;

Z is $-O-$, $-N(R^a)-$, $-S(O)_b-$ or $-CH(R^a)-$; wherein R^a is hydrogen or $C_{1-4}alkyl$ and b is 0-5 2;

R^8 is hydrogen, $C_{1-4}alkyl$, carbocyclyl or heterocyclyl; wherein R^8 may be optionally substituted on carbon by one or more substituents selected from R^{26} ; and wherein if said heterocyclyl contains an $-NH-$ group, that nitrogen may be optionally substituted by a group selected from R^{27} ;

10 R^9 is hydrogen or $C_{1-4}alkyl$;

R^{10} and R^{11} are independently selected from hydrogen, $C_{1-4}alkyl$, carbocyclyl or heterocyclyl; or R^{10} and R^{11} together form $C_{2-6}alkylene$; wherein R^{10} and R^{11} or R^{10} and R^{11} together may be independently optionally substituted on carbon by one or more substituents selected from R^{28} ; and wherein if said heterocyclyl contains an $-NH-$ moiety, that nitrogen may
15 be optionally substituted by one or more R^{29} ;


R^{12} is hydrogen, $C_{1-4}alkyl$, carbocyclyl or heterocyclyl; wherein R^{12} may be optionally substituted on carbon by one or more substituents selected from R^{30} ; and wherein if said heterocyclyl contains an $-NH-$ moiety, that nitrogen may be optionally substituted by one or more R^{31} ;

20 R^{13} is hydrogen, halo, nitro, cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, hydroxyaminocarbonyl, $C_{1-10}alkyl$, $C_{2-10}alkenyl$, $C_{2-10}alkynyl$, $C_{1-10}alkoxy$, $C_{1-10}alkoxycarbonyl$, $C_{1-10}alkanoyl$, $C_{1-10}alkanoyloxy$, $N-(C_{1-10}alkyl)amino$, $N,N-(C_{1-10}alkyl)_2amino$, $N,N,N-(C_{1-10}alkyl)_3ammonio$, $C_{1-10}alkanoylamino$, $N-(C_{1-10}alkyl)carbamoyl$, $N,N-(C_{1-10}alkyl)_2carbamoyl$, $C_{1-10}alkylS(O)_a$ wherein a is 0 to 2, $N-(C_{1-10}alkyl)sulphamoyl$,

25 $N,N-(C_{1-10}alkyl)_2sulphamoyl$, $N-(C_{1-10}alkyl)sulphamoylamino$,

$N,N-(C_{1-10}alkyl)_2sulphamoylamino$, $C_{1-10}alkoxycarbonylamino$, carbocyclyl, carbocyclyl $C_{1-10}alkyl$, heterocyclic group, heterocyclyl $C_{1-10}alkyl$, carbocyclyl- $(C_{1-10}alkylene)_e-R^{32}-(C_{1-10}alkylene)_f$ or


heterocyclyl- $(C_{1-10}alkylene)_g-R^{33}-(C_{1-10}alkylene)_h$; wherein R^{13} may be optionally substituted
30 on carbon by one or more substituents selected from R^{36} ; and wherein if said heterocyclyl contains an $-NH-$ group, that nitrogen may be optionally substituted by a group selected from R^{37} ; or R^{13} is a group of formula (IB):



(IB)

wherein:

- X is $-\text{N}(\text{R}^{38})-$, $-\text{N}(\text{R}^{38})\text{C}(\text{O})-$, $-\text{O}-$, and $-\text{S}(\text{O})_a-$; wherein a is 0-2 and R^{38} is hydrogen or
- 5 C_{1-4} alkyl;
- R^{14} is hydrogen or C_{1-4} alkyl;
- R^{15} and R^{16} are independently selected from hydrogen, halo, nitro, cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino,
- 10 N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkyl $\text{S}(\text{O})_a$ wherein a is 0 to 2, C_{1-6} alkoxycarbonyl, N -(C_{1-6} alkyl)sulphamoyl, N,N -(C_{1-6} alkyl) $_2$ sulphamoyl, carbocyclyl or heterocyclic group; wherein R^{15} and R^{16} may be independently optionally substituted on carbon by one or more substituents selected from R^{41} ; and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally substituted by a group selected from R^{42} ;
- 15 R^{17} is selected from hydrogen, halo, nitro, cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, hydroxyaminocarbonyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy, C_{1-10} alkanoyl, C_{1-10} alkanoyloxy, N -(C_{1-10} alkyl)amino, N,N -(C_{1-10} alkyl) $_2$ amino, C_{1-10} alkanoylamino, N -(C_{1-10} alkyl)carbamoyl, C_{1-10} alkoxycarbonyl, N,N -(C_{1-10} alkyl) $_2$ carbamoyl, C_{1-10} alkyl $\text{S}(\text{O})_a$ wherein a is 0 to 2, N -(C_{1-10} alkyl)sulphamoyl, N,N -(C_{1-10} alkyl) $_2$ sulphamoyl,
- 20 N -(C_{1-10} alkyl)sulphamoylamino, N,N -(C_{1-10} alkyl) $_2$ sulphamoylamino, carbocyclyl, carbocyclyl C_{1-10} alkyl, heterocyclic group, heterocyclyl C_{1-10} alkyl, carbocyclyl-(C_{1-10} alkylene) $_e$ - R^{43} -(C_{1-10} alkylene) $_f$ or heterocyclyl-(C_{1-10} alkylene) $_g$ - R^{44} -(C_{1-10} alkylene) $_h$; wherein R^{17} may be optionally substituted on carbon by one or more substituents selected from R^{47} ; and wherein if said heterocyclyl
- 25 contains an -NH- group, that nitrogen may be optionally substituted by a group selected from R^{48} ; or R^{17} is a group of formula (IC):



(IC)

wherein:

R^{18} is selected from hydrogen or C_{1-4} alkyl;

R^{19} is selected from hydrogen, halo, nitro, cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl,

- 5 C_{1-6} alkanoyloxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkylS(O) $_a$ wherein a is 0 to 2, C_{1-6} alkoxycarbonyl, N -(C_{1-6} alkyl)sulphamoyl, N,N -(C_{1-6} alkyl) $_2$ sulphamoyl, carbocyclyl or heterocyclic group; where R^{19} may be independently optionally substituted on carbon by one or more substituents selected from R^{51} ; and wherein if said heterocyclyl contains an -NH- group,
- 10 that nitrogen may be optionally substituted by a group selected from R^{52} ;

R^{20} is selected from halo, nitro, cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, hydroxyaminocarbonyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy, C_{1-10} alkoxycarbonyl, C_{1-10} alkanoyl, C_{1-10} alkanoyloxy, N -(C_{1-10} alkyl)amino, N,N -(C_{1-10} alkyl) $_2$ amino, N,N,N -(C_{1-10} alkyl) $_3$ ammonio, C_{1-10} alkanoylamino,

- 15 N -(C_{1-10} alkyl)carbamoyl, N,N -(C_{1-10} alkyl) $_2$ carbamoyl, C_{1-10} alkylS(O) $_a$ wherein a is 0 to 2, N -(C_{1-10} alkyl)sulphamoyl, N,N -(C_{1-10} alkyl) $_2$ sulphamoyl, N -(C_{1-10} alkyl)sulphamoylamino, N,N -(C_{1-10} alkyl) $_2$ sulphamoylamino, C_{1-10} alkoxycarbonylamino, carbocyclyl, carbocyclyl C_{1-10} alkyl, heterocyclic group, heterocyclyl C_{1-10} alkyl, carbocyclyl-(C_{1-10} alkylene) $_e$ - R^{53} -(C_{1-10} alkylene) $_f$ or
- 20 heterocyclyl-(C_{1-10} alkylene) $_g$ - R^{54} -(C_{1-10} alkylene) $_h$; wherein R^{20} may be independently optionally substituted on carbon by one or more R^{57} ; and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally substituted by a group selected from R^{58} ;

p is 1-3; wherein the values of R^{15} may be the same or different;

q is 0-1;

- 25 r is 0-3; wherein the values of R^{16} may be the same or different;

m is 0-2; wherein the values of R^{12} may be the same or different;

n is 1-2; wherein the values of R^8 may be the same or different;

z is 0-3; wherein the values of R^{19} may be the same or different;

R^{21} is selected from hydrogen or C_{1-6} alkyl;

- 30 R^{22} and R^{23} are independently selected from hydrogen, hydroxy, amino, mercapto, C_{1-6} alkyl, C_{1-6} alkoxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkylS(O) $_a$ wherein a is 0 to 2;

R^{24} is selected from hydrogen, hydroxy, C_{1-6} alkyl, C_{1-4} alkoxy and C_{1-6} alkanoyloxy;

R^{25} is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, C_{1-4} alkanoyl, C_{1-4} alkanoyloxy, N -(C_{1-4} alkyl)amino, N,N -(C_{1-4} alkyl)₂amino, C_{1-4} alkanoylamino, N -(C_{1-4} alkyl)carbamoyl, 5 N,N -(C_{1-4} alkyl)₂carbamoyl, C_{1-4} alkylS(O)_a wherein a is 0 to 2, C_{1-4} alkoxycarbonyl, N -(C_{1-4} alkyl)sulphamoyl and N,N -(C_{1-4} alkyl)₂sulphamoyl; wherein R^{25} , may be independently optionally substituted on carbon by one or more R^{67} ;

R^{26} , R^{28} , R^{30} , R^{36} , R^{41} , R^{47} , R^{51} and R^{57} are independently selected from halo, nitro, cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, hydroxyaminocarbonyl, C_{1-10} alkyl, 10 C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy, C_{1-10} alkanoyl, C_{1-10} alkanoyloxy, C_{1-10} alkoxycarbonyl, N -(C_{1-10} alkyl)amino, N,N -(C_{1-10} alkyl)₂amino, N,N,N -(C_{1-10} alkyl)₃ammonio, C_{1-10} alkanoylamino, N -(C_{1-10} alkyl)carbamoyl, N,N -(C_{1-10} alkyl)₂carbamoyl, C_{1-10} alkylS(O)_a wherein a is 0 to 2, N -(C_{1-10} alkyl)sulphamoyl, N,N -(C_{1-10} alkyl)₂sulphamoyl, N -(C_{1-10} alkyl)sulphamoylamino, N,N -(C_{1-10} alkyl)₂sulphamoylamino, C_{1-10} alkoxycarbonylamino, carbocyclyl, 15 carbocyclyl C_{1-10} alkyl, heterocyclic group, heterocyclyl C_{1-10} alkyl, carbocyclyl-(C_{1-10} alkylene)_e- R^{59} -(C_{1-10} alkylene)_i- or heterocyclyl-(C_{1-10} alkylene)_g- R^{60} -(C_{1-10} alkylene)_h-; wherein R^{26} , R^{28} , R^{30} , R^{36} , R^{41} , R^{47} , R^{51} and R^{57} may be independently optionally substituted on carbon by one or more R^{63} ; and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally substituted by a 20 group selected from R^{64} ;

R^{27} , R^{29} , R^{31} , R^{37} , R^{42} , R^{48} , R^{52} , R^{58} and R^{64} are independently selected from C_{1-6} alkyl, C_{1-6} alkanoyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -(C_{1-6} alkyl)sulphamoyl, N,N -(C_{1-6} alkyl)₂sulphamoyl, C_{1-6} alkoxycarbonyl, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl)₂carbamoyl, benzyl, phenethyl, benzoyl, phenylsulphonyl and phenyl;

25 R^{32} , R^{33} , R^{43} , R^{44} , R^{53} , R^{54} , R^{59} and R^{60} are independently selected from -O-, - NR^{65} -, -S(O)_x-, - $NR^{65}C(O)NR^{66}$ -, - $NR^{65}C(S)NR^{66}$ -, -OC(O)N=C-, - $NR^{65}C(O)$ - or -C(O) NR^{65} -; wherein R^{65} and R^{66} are independently selected from hydrogen or C_{1-6} alkyl, and x is 0-2;

R^{63} and R^{67} are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, methyl, 30 ethyl, methoxy, ethoxy, vinyl, allyl, ethynyl, methoxycarbonyl, formyl, acetyl, formamido, acetylamino, acetoxymethyl, methylamino, dimethylamino, N -methylcarbamoyl,

N,N-dimethylcarbamoyl, methylthio, methylsulphinyl, mesyl, *N*-methylsulphamoyl and *N,N*-dimethylsulphamoyl; and


e, f, g and h are independently selected from 0-2;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

A B S T R A C T

Title: CHEMICAL COMPOUNDS

Compounds of formula (I):



5

(I)

pharmaceutically acceptable salts, solvates, solvates of such salts and prodrugs thereof and their use as ileal bile acid transport (IBAT) inhibitors for the treatment of hyperlipidaemia are described. Processes for their manufacture and pharmaceutical compositions containing them

10 are also described.